



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>

Tolman, Cyrus Fisher, 1873-

553.16
T652


A study of the magmatic sulfid ores.

553.16
T652





**BRANNER
GEOLOGICAL LIBRARY**



Gift of

MRS. A. P. BRANNER

*Dr. J. F. Newsom
with the regards of the authors*

LELAND STANFORD JUNIOR UNIVERSITY PUBLICATIONS
UNIVERSITY SERIES

A Study
of the
Magmatic Sulfid Ores

BY

C. F. TOLMAN, JR.

Associate Professor of Economic Geology

and

AUSTIN F. ROGERS

Associate Professor of Mineralogy and Petrography

With 20 Plates and 7 Text Figures

STANFORD UNIVERSITY, CALIFORNIA
PUBLISHED BY THE UNIVERSITY
1916

UNIVERSITY SERIES

- INHERITANCE IN SILKWORMS, I.** Vernon Lyman Kellogg, Professor of Entomology. 89 pp., 4 plates. 1908. Price \$1.00.
- THE OPISTHOBRANCHIATE MOLLUSCA OF THE BRANNER-AGASSIZ EXPEDITION TO BRAZIL.** Frank Mace McFarland, Professor of Histology. 105 pp., 19 plates. 1909. Price, \$1.00.
- A STUDY OF THE NORMAL CONSTITUENTS OF THE POTABLE WATER OF THE SAN FRANCISCO PENINSULA.** John Pearce Mitchell, Assistant Professor of Chemistry. 70 pp., 1 map. 1910. Price, 50c.
- SYNOPSIS OF THE TRUE CRABS (BRACHYURA) OF MONTEREY BAY, CALIFORNIA.** Frank Walter Weymouth. 64 pp., 14 plates. 1910. Price, 50c.
- THE OSTEOLOGY OF CERTAIN SCOMBROID FISHES.** Edwin Chapin Starks, Assistant Professor of Zoology. 49 pp., 2 plates, 1 text figure. 1911. Price, 50c.
- A PHYSICAL THEORY OF ELECTRIFICATION.** Fernando Sanford, Professor of Physics. 69 pp., 2 plates. 1911. Price, 50c.
- THE MATZKE MEMORIAL VOLUME.** Papers by John Ernst Matzke, late Professor of Romanic Languages, and Thirteen Colleagues. 162 pp. 1911. Price, \$1.00.
- DAS HISTORISCHE PRÄSENS IN DER ÄLTEREN DEUTSCHEN SPRACHE.** Bruno Boezinger, Assistant Professor of Germanic Languages. 91 pp. 1912. Price, 50c.
- THE EFFECT OF A STRICTLY VEGETABLE DIET ON THE SPONTANEOUS ACTIVITY, THE RATE OF GROWTH, AND THE LONGEVITY OF THE ALBINO RAT.** James Rollin Slonaker, Assistant Professor of Physiology. 36 pp., 1 plate, 15 text figures. 1912. Price, 50c.
- CATALOGUE DE TOUS LES LIVRES DE FEU M. CHAPELAIN.** (Bibliothèque Nationale, Fonds Français, Nouv. Acq., No. 318.) Colbert Searles, Associate Professor of Romanic Languages. 119 pp., 2 plates. 1912. Price, 75c.
- THE DUDLEY MEMORIAL VOLUME.** Papers by William Russel Dudley, late Professor of Botany, and Several Colleagues. 137 pp., 12 text figures, 9 plates. 1913. Price, \$1.00.
- THE FISHES OF THE STANFORD EXPEDITION TO BRAZIL.** Edwin Chapin Starks, Assistant Professor of Zoology. 77 pp., 15 plates. 1913. Price, 75c.
- THE BIRDS OF THE LATIN POETS.** Ernest Whitney Martin, Associate Professor of Greek. 260 pp. 1914. Price, \$1.00.

(Continued on third page of cover.)

UNIVERSITY SERIES

- INHERITANCE IN SILKWORMS, I.** Vernon Lyman Kellogg, Professor of Entomology. 89 pp., 4 plates. 1908. Price \$1.00.
- THE OPISTHOBRANCHIATE MOLLUSCA OF THE BRANNER-AGASSIZ EXPEDITION TO BRAZIL.** Frank Mace McFarland, Professor of Histology. 105 pp., 19 plates. 1909. Price, \$1.00.
- A STUDY OF THE NORMAL CONSTITUENTS OF THE POTABLE WATER OF THE SAN FRANCISCO PENINSULA.** John Pearce Mitchell, Assistant Professor of Chemistry. 70 pp., 1 map. 1910. Price, 50c.
- SYNOPSIS OF THE TRUE CRABS (BRACHYURA) OF MONTEREY BAY, CALIFORNIA.** Frank Walter Weymouth. 64 pp., 14 plates. 1910. Price, 50c.
- THE OSTEOLOGY OF CERTAIN SCOMBROID FISHES.** Edwin Chapin Starks, Assistant Professor of Zoology. 49 pp., 2 plates, 1 text figure. 1911. Price, 50c.
- A PHYSICAL THEORY OF ELECTRIFICATION.** Fernando Sanford, Professor of Physics. 69 pp., 2 plates. 1911. Price, 50c.
- THE MATZKE MEMORIAL VOLUME.** Papers by John Ernst Matzke, late Professor of Romanic Languages, and Thirteen Colleagues. 162 pp. 1911. Price, \$1.00.
- DAS HISTORISCHE PRÄSENS IN DER ÄLTEREN DEUTSCHEN SPRACHE.** Bruno Boezinger, Assistant Professor of Germanic Languages. 91 pp. 1912. Price, 50c.
- THE EFFECT OF A STRICTLY VEGETABLE DIET ON THE SPONTANEOUS ACTIVITY, THE RATE OF GROWTH, AND THE LONGEVITY OF THE ALBINO RAT.** James Rollin Slonaker, Assistant Professor of Physiology. 36 pp., 1 plate, 15 text figures. 1912. Price, 50c.
- CATALOGUE DE TOUS LES LIVRES DE FEU M. CHAPELAIN.** (Bibliothèque Nationale, Fonds Français, Nouv. Acq., No. 318.) Colbert Searles, Associate Professor of Romanic Languages. 119 pp., 2 plates. 1912. Price, 75c.
- THE DUDLEY MEMORIAL VOLUME.** Papers by William Russel Dudley, late Professor of Botany, and Several Colleagues. 137 pp., 12 text figures, 9 plates. 1913. Price, \$1.00.
- THE FISHES OF THE STANFORD EXPEDITION TO BRAZIL.** Edwin Chapin Starks, Assistant Professor of Zoology. 77 pp., 15 plates. 1913. Price, 75c.
- THE BIRDS OF THE LATIN POETS.** Ernest Whitney Martin, Associate Professor of Greek. 260 pp. 1914. Price, \$1.00.

(Continued on third page of cover.)



LELAND STANFORD JUNIOR UNIVERSITY PUBLICATIONS
UNIVERSITY SERIES

A Study
of the
Magmatic Sulfid Ores

BY

C. F. TOLMAN, JR.

Associate Professor of Economic Geology

and

AUSTIN F. ROGERS

Associate Professor of Mineralogy and Petrography

With 20 Plates and 7 Text Figures

STANFORD LIBRARY

STANFORD UNIVERSITY, CALIFORNIA

PUBLISHED BY THE UNIVERSITY

1916

11

553.16
T652

STANFORD UNIVERSITY
PRESS

631746

YNA334 : 631746

CONTENTS		PAGE
INTRODUCTION		5
Definition of magmatic ores; Theories as to the segregation of magmatic ores; Classification of magmatic ores; Deposits studied; Plan of treatment.		
PART I. DISCUSSION OF THE BEARING OF MAGMATIC DIFFERENTIATION ON ORE SEGREGATION		9
Review of the current theories of magmatic differentiation; Theories favored by us; Problems to be investigated; Methods of investigation; Microscopic investigation; Conclusions from our microscopic investigations; The evidence on which the above conclusions are based; Replacement phenomena in the magmatic ores; Alteration and later rearrangement in magmatic ores; Absence of pneumatolytic and hydrothermal alteration during ore deposition; Magnetite-ilmenite magmatic ores; Euhedral magnetite formed at a late magmatic stage; Other accessory minerals of probable late magmatic origin; Stages recognized in the formation of magmatic ores.		
PART II. DESCRIPTION OF THE VARIOUS DEPOSITS OF THE MAGMATIC ORES		
GROUP I. THE NICKEL- AND COPPER-BEARING PYRRHOTITIC DEPOSITS		
SUDBURY, CANADA		23
Geology; Structure of the ore bodies; Microscopic descriptions; Origin of the Sudbury ores; Bibliography of the Sudbury ore deposits.		
THE ALEXO MINE, ONTARIO, CANADA		37
Geology; Uglow's conclusions; Microscopic description; Summary; Bibliography of the Alexo ore deposits.		
THE FRIDAY MINE, SAN DIEGO COUNTY, CALIFORNIA		39
Microscopic description; Summary.		
THE GOLDEN CURRY MINE, MONTANA		41
Geology; Microscopic description.		
PROSPECT HILL, LITCHFIELD, CONNECTICUT		43
KNOX COUNTY, MAINE		44
MOUNTAIN, WISCONSIN		44
OTHER PYRRHOTITE DEPOSITS IN THE UNITED STATES		44
INSIZWA RANGE, EAST GRICUALAND, SOUTH AFRICA		44
NORWAY		46
Geology; Petrography and mineralogy; Discussion of Vogt's conclusions; Summary; Bibliography of the Norwegian deposits.		

	PAGE
OTHER EUROPEAN OCCURRENCES	54
Sweden; Baden, Horbach, and Todtmoos, Germany; Sohland and Sweiderich on the boundary between Saxony and Bohemia.	
GROUP II. THE MAGMATIC CHALCOPYRITE-BORNITE DEPOSITS	
OOKIEP, NAMAQUALAND, SOUTH AFRICA	56
Geology; Microscopic descriptions; Summary; Bibliography of the Ookiep deposits.	
ENGELS MINE, PLUMAS COUNTY, CALIFORNIA	61
Geology; Microscopic descriptions; Summary; Bibliography of the Engels ore deposit.	
REMARKS ON CERTAIN OTHER DEPOSITS THAT HAVE BEEN CLASSIFIED AS MAGMATIC	
PYRITIC DEPOSITS	65
MAGNETITE-ILMENITE DEPOSITS	67
OTHER MAGMATIC IRON ORES	67
CHROMITE DEPOSITS	68
PART III. SUMMARY AND CONCLUSIONS	
CRITERIA FOR THE RECOGNITION OF MAGMATIC ORES	69
SUMMARY OF THE CHARACTERISTICS OF MAGMATIC ORES	70
<hr/>	
EXPLANATION OF PLATES, AND METHODS OF PREPARING PHOTOGRAPHS AND SECTIONS	74
PLATES, I-XX	

ACKNOWLEDGMENTS

This study has been made possible by the generous contribution of material by the following named gentlemen: F. L. Hess, Ernest Howe, E. P. Jennings, J. F. Kemp, C. W. Knight, Adolph Knopf, R. D. Longyear, F. H. Mason, M. E. Morgan, Heinrich Ries, A. W. Rogers, Beecher Sterne, H. W. Turner, and T. L. Walker.

We also acknowledge our indebtedness to Professor Bailey Willis for a critical reading of the manuscript and for valuable suggestions.

A STUDY OF THE MAGMATIC SULFID ORES

INTRODUCTION

DEFINITION OF MAGMATIC ORES

The term "magmatic ore" is generally applied to those phases of igneous rocks in which there has been an unusual accumulation, supposedly during the molten stage, of the accessory ore-minerals. The recognition of this type of ore deposits is due largely to the work of Vogt, who, in a classic series of papers,¹ not only established the existence of magmatic ores, but also arranged them into well defined groups, and gave the characteristics of each group. His classification has been followed generally by the authors of recent textbooks.² Nevertheless some confusion is apparent in geological literature as to the meaning of the term "magmatic" as applied to ore deposits. This is due in part to the designation³ of all deposits of direct or indirect magmatic origin as magmatic deposits. For example, few doubt the magmatic origin of contact deposits and of cassiterite-tourmaline-quartz veins; but these are not magmatic deposits or segregations in the strict sense of the term.

The term "magmatic deposits" should be limited to those segregations of ore-minerals that take place under the influence of, or closely connected with, the molten stage of the parent rock. Ore accumulations accompanied by destructive pneumatolytic action, or those formed by hydrothermal solutions, are not to be classed as magmatic deposits, altho they may be closely related to, and follow, the magmatic period

¹ Vogt, J. H. L.—Bildung von Erzlagerstätten durch Differentiationsprocesse in basischen Eruptivmagmata. Zeit. f. prakt. Geol., Jahrgang 1893, 4-11, 125-43, 257-284.

—Beiträge zur genetischen Classification der durch magmatische Differentiation-processe und der durch Pneumatolyse entstanden Erzvorkommen. Zeit. f. prakt. Geol., Jahrg. 1894, 381-399.

—Weitere Untersuchungen über die Ausscheidungen von Titaneisenerzen in basischen Eruptivgesteine. Zeit. f. prakt. Geol., Jahrg. 1900, 233-242, 370-382; Jahrg. 1901, 9-19, 180-186, 289-296, 327-340.

² As is noted below, Vogt classifies as magmatic certain pyritic deposits that we believe should not be included in the magmatic group, and also omits one of the most important divisions of this group, viz., the chalcopyrite-bornite type of magmatic ores.

³ Geijer, Per.—Iron-ore geology in Sweden and America. Econ. Geol., 10, 231 (1915).

of ore concentration. In as much as ore concentrations connected with persilicic ("acid") rocks are of the latter type, the typical magmatic deposits are confined to the subsilicic ("basic") rocks.

THEORIES AS TO THE SEGREGATION OF MAGMATIC ORES

Altho the chief types of magmatic ores are well recognized, there is considerable difference of opinion as to the details of the processes by which the ores are formed, and as to the "order of crystallization" of the ore-minerals with reference to the silicates. These different ideas may be grouped broadly as follows:

1. Magmatic ores have been defined as unusual accumulations of certain of the accessory minerals of the igneous rocks; and as the accessory minerals are generally believed to be the first to crystallize out of the magma, according to the order of crystallization suggested by Rosenbusch, the natural inference is that the ores are the first to form, and that they settle by gravity to the base of the magma. For example, it is generally assumed that the segregation of the large bodies of iron ore in the Bushveldt laccolith in South Africa and in the Duluth laccolith and the sulfid masses in the Sudbury laccolith have been controlled by gravity.⁴

Beyschlag, Krusch, and Vogt⁵ state "Bei den meisten Eruptivgesteinen beginnt die Kristallisation mit der Aussonderung der sogenannten Erzminerale, wie Magnetit oder Titanomagnetit, Eisenglanz, Ilmenit, Zirkon, Apatit, Schwefelkies, mitunter auch Spinel u. s. w. In einer etwas späteren Stufe der Verfestigung beginnen die Eisenmagnesiumsilikate, wie Glimmer, Hornblend, und Pyroxenminerale und Olivin, zu kristallisieren."

2. Vogt, who has examined the magmatic ores both in the field and with the microscope, favors the hypothesis that prior to the crystallization of the rock-minerals the ores separate as an immiscible sulfid or oxid melt, which continues in the molten state during the consolidation of the silicates, intrudes the latter, and finally crystallizes.

3. Modern investigation of polished surfaces of ores has led to the discovery that the magmatic sulfid ore-minerals have a definite order of crystallization, and that the younger minerals intrude and replace the older. To meet this, Howe has framed a modification of Vogt's hypothesis. He suggests⁶ that the different minerals which make up the sulfid

⁴ Daly, R. A.—Igneous rocks and their origin, 454 (1914).

⁵ Die Erzlagerstätten der Nutzbaren Mineralien und Gesteine, 1, 240 (1910).

⁶ Howe, E.—Petrographical notes on the Sudbury nickel deposits. Econ. Geol., 9, 522 (1914).

matte are mutually immiscible, and the sulfids last to crystallize intrude the previously solidified sulfids.

4. Our study of the magmatic ores has led us to frame the hypothesis that *the magmatic ores in general have been introduced at a late magmatic stage as a result of mineralizers, and that the ore-minerals replace the silicates. This replacement, however, differs from that caused by destructive pneumatolytic or hydrothermal processes in that quartz and secondary silicates are not formed at the time the ores are deposited.*

None of the magmatic ores are entirely free from alteration, and in some cases they are associated with high-temperature alteration products as well as with those of hydrothermal origin. It became necessary for us, therefore, to study the various stages of mineral deposition, especially the migrations and alterations that follow the original deposition of the ore.

Some of the data indicating that the magmatic ores are later than the silicates have influenced a number of geologists to question the propriety of classifying certain deposits as magmatic and perhaps to doubt the importance of the type as a whole. Our work meets the objections urged by opponents of the magmatic theories in that it proves the ores are formed within the magmatic period, as defined by us, altho the ore-minerals are later than the silicates.


CLASSIFICATION OF THE MAGMATIC ORES

Vogt[†] separates the magmatic ores according to mineral composition into three groups, viz.: the oxid ores, the sulfid ores, and those of the native metals. The last group includes nickeliferous iron, platinum, copper, and gold. The magmatic deposits of nickeliferous iron and of platinum are curiosities rather than ore deposits. The existence of magmatic copper and gold has been asserted, but not proved by accurate microscopic work. Therefore they are not considered further by us. The oxid group includes deposits of chromite, corundum, and, most important, of magnetite and ilmenite. Altho we touch upon these briefly, the study presented here deals chiefly with the magmatic sulfid deposits. These sulfid ores have been classified as magmatic because of their exclusive occurrence in igneous rocks; because the ore formation is not accompanied by pneumatolytic, contact, or hydrothermal silicates, so markedly developed in connection with sulfid deposits of the non-magmatic types in igneous rocks; and because the

[†] Zeit. f. prakt. Geol., Jahrgang 1894, p. 382.



**BRANNER
GEOLOGICAL LIBRARY**



Gift of

Mrs. A. P. NELSON

*Dr. J. F. Newcom
with the regards of the authors.*

LELAND STANFORD JUNIOR UNIVERSITY PUBLICATIONS
UNIVERSITY SERIES

A Study
of the
Magmatic Sulfid Ores

BY

C. F. TOLMAN, JR.

Associate Professor of Economic Geology

and

AUSTIN F. ROGERS

Associate Professor of Mineralogy and Petrography

With 20 Plates and 7 Text Figures

STANFORD UNIVERSITY, CALIFORNIA
PUBLISHED BY THE UNIVERSITY
1916

UNIVERSITY SERIES

- INHERITANCE IN SILKWORMS, I.** Vernon Lyman Kellogg, Professor of Entomology. 89 pp., 4 plates. 1908. Price \$1.00.
- THE OPISTHOBRANCHIATE MOLLUSCA OF THE BRANNER-AGASSIZ EXPEDITION TO BRAZIL.** Frank Mace McFarland, Professor of Histology. 105 pp., 19 plates. 1909. Price, \$1.00.
- A STUDY OF THE NORMAL CONSTITUENTS OF THE POTABLE WATER OF THE SAN FRANCISCO PENINSULA.** John Pearce Mitchell, Assistant Professor of Chemistry. 70 pp., 1 map. 1910. Price, 50c.
- SYNOPSIS OF THE TRUE CRABS (BRACHYURA) OF MONTEREY BAY, CALIFORNIA.** Frank Walter Weymouth. 64 pp., 14 plates. 1910. Price, 50c.
- THE OSTEOLOGY OF CERTAIN SCOMBROID FISHES.** Edwin Chapin Starks, Assistant Professor of Zoology. 49 pp., 2 plates, 1 text figure. 1911. Price, 50c.
- A PHYSICAL THEORY OF ELECTRIFICATION.** Fernando Sanford, Professor of Physics. 69 pp., 2 plates. 1911. Price, 50c.
- THE MATZKE MEMORIAL VOLUME.** Papers by John Ernst Matzke, late Professor of Romanic Languages, and Thirteen Colleagues. 162 pp. 1911. Price, \$1.00.
- DAS HISTORISCHE PRÄSENS IN DER ÄLTEREN DEUTSCHEN SPRACHE.** Bruno Boezinger, Assistant Professor of Germanic Languages. 91 pp. 1912. Price, 50c.
- THE EFFECT OF A STRICTLY VEGETABLE DIET ON THE SPONTANEOUS ACTIVITY, THE RATE OF GROWTH, AND THE LONGEVITY OF THE ALBINO RAT.** James Rollin Slonaker, Assistant Professor of Physiology. 36 pp., 1 plate, 15 text figures. 1912. Price, 50c.
- CATALOGUE DE TOUS LES LIVRES DE FEU M. CHAPELAIN.** (Bibliothèque Nationale, Fonds Français, Nouv. Acq., No. 318.) Colbert Searles, Associate Professor of Romanic Languages. 119 pp., 2 plates. 1912. Price, 75c.
- THE DUDLEY MEMORIAL VOLUME.** Papers by William Russel Dudley, late Professor of Botany, and Several Colleagues. 137 pp., 12 text figures, 9 plates. 1913. Price, \$1.00.
- THE FISHES OF THE STANFORD EXPEDITION TO BRAZIL.** Edwin Chapin Starks, Assistant Professor of Zoology. 77 pp., 15 plates. 1913. Price, 75c.
- THE BIRDS OF THE LATIN POETS.** Ernest Whitney Martin, Associate Professor of Greek. 260 pp. 1914. Price, \$1.00.

(Continued on third page of cover.)

UNIVERSITY SERIES

- INHERITANCE IN SILKWORMS, I. Vernon Lyman Kellogg, Professor of Entomology. 89 pp., 4 plates. 1908. Price \$1.00.
- THE OPISTHOBRANCHIATE MOLLUSCA OF THE BRANNER-AGASSIZ EXPEDITION TO BRAZIL. Frank Mace McFarland, Professor of Histology. 105 pp., 19 plates. 1909. Price, \$1.00.
- A STUDY OF THE NORMAL CONSTITUENTS OF THE POTABLE WATER OF THE SAN FRANCISCO PENINSULA. John Pearce Mitchell, Assistant Professor of Chemistry. 70 pp., 1 map. 1910. Price, 50c.
- SYNOPSIS OF THE TRUE CRABS (BRACHYURA) OF MONTEREY BAY, CALIFORNIA. Frank Walter Weymouth. 64 pp., 14 plates. 1910. Price, 50c.
- THE OSTEOLOGY OF CERTAIN SCOMBROID FISHES. Edwin Chapin Starks, Assistant Professor of Zoology. 49 pp., 2 plates, 1 text figure. 1911. Price, 50c.
- A PHYSICAL THEORY OF ELECTRIFICATION. Fernando Sanford, Professor of Physics. 69 pp., 2 plates. 1911. Price, 50c.
- THE MATZKE MEMORIAL VOLUME. Papers by John Ernst Matzke, late Professor of Romanic Languages, and Thirteen Colleagues. 162 pp. 1911. Price, \$1.00.
- DAS HISTORISCHE PRÄSENS IN DER ÄLTEREN DEUTSCHEN SPRACHE. Bruno Boezinger, Assistant Professor of Germanic Languages. 91 pp. 1912. Price, 50c.
- THE EFFECT OF A STRICTLY VEGETABLE DIET ON THE SPONTANEOUS ACTIVITY, THE RATE OF GROWTH, AND THE LONGEVITY OF THE ALBINO RAT. James Rollin Slonaker, Assistant Professor of Physiology. 36 pp., 1 plate, 15 text figures. 1912. Price, 50c.
- CATALOGUE DE TOUS LES LIVRES DE FEU M. CHAPELAIN. (Bibliothèque Nationale, Fonds Français, Nouv. Acq., No. 318.) Colbert Searles, Associate Professor of Romanic Languages. 119 pp., 2 plates. 1912. Price, 75c.
- THE DUDLEY MEMORIAL VOLUME. Papers by William Russel Dudley, late Professor of Botany, and Several Colleagues. 137 pp., 12 text figures, 9 plates. 1913. Price, \$1.00.
- THE FISHES OF THE STANFORD EXPEDITION TO BRAZIL. Edwin Chapin Starks, Assistant Professor of Zoology. 77 pp., 15 plates. 1913. Price, 75c.
- THE BIRDS OF THE LATIN POETS. Ernest Whitney Martin, Associate Professor of Greek. 260 pp. 1914. Price, \$1.00.

(Continued on third page of cover.)





LELAND STANFORD JUNIOR UNIVERSITY PUBLICATIONS
UNIVERSITY SERIES

A Study
of the
Magmatic Sulfid Ores

BY

C. F. TOLMAN, JR.

Associate Professor of Economic Geology

and

AUSTIN F. ROGERS

Associate Professor of Mineralogy and Petrography

With 20 Plates and 7 Text Figures

STANFORD LIBRARY

STANFORD UNIVERSITY, CALIFORNIA

PUBLISHED BY THE UNIVERSITY

1916

16

553.16
T652

STANFORD UNIVERSITY
PRESS

631746

YPA: 1907A73

of secondary migration of ore-minerals, even where they are cut by veinlets of chlorite. The striking thing about the magmatic ores in general is the slight amount of rearrangement they have undergone.

The results of our study of the ores of the Alexo mine, Canada, are of interest in showing that, altho the silicates are intensely altered, and completely serpentinized, nevertheless the ore is only slightly affected. Here the serpentinization is subsequent to ore formation, as shown by veinlets of serpentine cutting the main ore masses. Two generations of serpentinization are recognized; the first accomplished the segregation of insignificant veinlets and specks of chalcopyrite within the area serpentinized, but apparently did not modify even the outline of main masses of ore; the second generation of serpentine is accompanied by still more minute scattered microscopic specks of ore. Only a small portion of the total ore suffered rearrangement during the process of serpentinization.

ABSENCE OF PNEUMATOLYTIC AND HYDROTHERMAL ALTERATION DURING ORE DEPOSITION

One of the characteristics of magmatic ores that has long been emphasized in the literature is the lack of secondary silicates produced by pneumatolytic and hydrothermal processes. In a few cases, however, the ores are accompanied by high-temperature minerals, such as garnet and tourmaline, and show gradations, therefore, towards the groups of high-temperature epigenetic ore deposits. Our work shows that hydrothermal alteration is invariably later than the period of magmatic ore formation, and therefore emphasizes the lack of alteration during the formation of the ores. It has been argued that if mineralizers or solutions are involved in the formation of the magmatic ores, they must of necessity not only dissolve the rock and deposit the ore, but must also react with the rock-forming minerals to produce secondary silicates. For example, Vogt⁸¹ was at first inclined, from field and microscopic data, to attribute the origin of the nickeliferous sulfids of Norway to "pneumatolytic" action; but the lack of the products of destructive pneumatolysis evidently caused him to abandon the idea that mineralizers are the agents of ore concentration, and he adopted the concept of an intrusive sulfid magma.

The idea that alteration invariably accompanies ore deposition is probably due to a considerable extent to the emphasis given rock alter-

⁸¹ [Geol. Fören. Förh., 1883] cited by Beyschlag, Krusch, Vogt, Erzlagertätten, p. 285.

ation in the classic paper of Lindgren.⁸² As a matter of fact, a careful microscopic examination by us of many types of non-magmatic ores has discovered veinlets or portions of veinlets along which the rock is little altered; and where the rock is affected, this alteration may be earlier or later than the deposition of the sulfids, and not connected directly with it. In replacement deposits in limestone, the most easily altered of all rocks, the ore often lies in contact with the limestone without intervening secondary minerals. *A priori*, no reason can be found why the composition and temperature of the ore-forming solutions in general should not be such that solution of the rock-minerals and deposition of the ore-minerals may take place without the formation of secondary silicates. *A fortiori*, this would be expected in the case of the ores under discussion. In the late stages of the consolidation of igneous rocks, constructive action of mineralizers aids and controls the formation of minerals without the development of secondary silicates. We have much to learn from the French scientists of the importance of mineralizers⁸³ in the crystallization of igneous rocks.

Extensive destructive pneumatolysis often occurs in connection with certain stages in the formation of pegmatite dikes and high-temperature veins, and results in the development of such minerals as quartz, muscovite, tourmaline, topaz, scapolite, etc. The temperature and character of the solutions and mineralizers are such that they are not in equilibrium with the minerals attacked, and they contain chemically active gases, such as boron, chlorine, fluorine, etc. They therefore develop in the country rock the complicated set of silicates mentioned.

⁸² Lindgren, W.—Metasomatic processes in fissure-veins. Trans. Am. Inst. Min. Eng., 30, 578-692 (1900).

⁸³ There is a striking difference between the German and American schools of petrography and the views of the French scientists. The former have interpreted the conditions governing the crystallization of deep-seated magmas in the light of observations on surface lavas and experiments on anhydrous melts. The latter have been impressed with the fumarolic action around volcanic vents, and have therefore concluded that mineralizers are important components in intrusive magmas. [Lacroix, A.—Les minéraux des fumerolles de l'éruption du Vésuve en avril 1906. Bull. Fr. Soc. Min., 30, 219-266 (1907).] Extreme views in regard to the role of mineralizers have been expressed by de Launay [Traité de Metallogénie. Gîtes minéraux et métallifères, 1, chap. 1 (1913)] who has developed a theoretical concept of the constitution of the earth involving a central metallic core with a superficial slag or silicate crust. He believes that mineralizers react upon the "slag," producing granite and other rocks, and form various types of ore bodies from the metals they have "extracted" from the metallic core.

We conclude that the magmatic ores, in contrast with the pegmatites and "high-temperature veins," occur in rocks which are little affected by the destructive action accompanying the latter.

The lack of alteration during the formation of the magmatic ores in "basic" rocks is probably the result of chemical composition rather than of temperature. The mineralizers of the "basic" rocks containing the magmatic sulfids produce hornblendization and biotitization, while those of "acid" rocks produce marked destructive effects (greisenization, tourmalinization, silicification, contact action, etc.).

The lack of alteration during ore formation, the fact that ore formation is often followed by the intrusion of pegmatitic differentiates, also the fact that the ores are limited to the parent "basic" rock and migrate into the intruded rock only to a very minor degree, are the chief characteristics of this definite and recognizable type of ore which has been designated as magmatic. Therefore we may well retain this term, even though certain misconceptions have been attached to it, such as consolidation in the early molten stage, injection of molten sulfids, etc.; recognizing that the microscopic examination definitely proves only that the ores are later than the primary silicates and earlier than the hydrothermal silicates, and also that the period of metallization is further removed from the main stages of rock consolidation than is generally believed.^{22a}

MAGNETITE-ILMENITE MAGMATIC ORES

We have studied, chiefly, the magmatic sulfid ores, and for these there is no doubt that the ores are later than the silicates. The same is true for the magnetite-ilmenite ores we have examined; and a review of the literature of these deposits containing accurate microscopic descriptions shows this to be true for all the iron ores of this type, although the bearing of this fact on the theories of magmatic differentiation does not appear to have been duly emphasized or appreciated.

EUHEDRAL MAGNETITE FORMED AT A LATE MAGMATIC STAGE

Berg²⁴ recognizes in the majority of magmatic iron-oxid deposits two generations of magnetite: euhedral magnetite earlier than the sili-

^{22a} At various times during the progress of our studies, we have considered the advisability of discarding the term "magmatic ore" in favor of some such name as epimagmatic. The fact that we have not been able to discover ores that show evidence of having been formed during the early stages in the consolidation of the magma, that the activity of mineralizers increases as crystallization progresses, and that any division between earlier and later minerals would be arbitrary, has deterred us from suggesting any departure from the present nomenclature, although such

cates, and larger areas of magnetite later than, and "corroding," the silicates. As magnetite is also a constituent of the magmatic sulfid ores, and as it occurs both as euhedral crystals and anhedral masses replacing the silicates, we were able to investigate the question as to the occurrence of the two generations of magnetite. We find that there are all gradations between small euhedral crystals and the large irregular areas of magnetite. The points of the larger masses may show crystal faces where they penetrate well into the silicate minerals. The magnetite appears to develop irregular forms where it surrounds the silicates, apparently following the lines of least resistance along the mineral boundaries, but develops euhedral forms within the silicates, where no line of weakness disturbs its tendency to crystallize regularly. We believe that the magnetite in the magmatic sulfid ores, and probably in the magmatic ores in general, is all deposited later than the silicates.

OTHER ACCESSORY MINERALS OF PROBABLE LATE MAGMATIC ORIGIN

The accessory minerals of igneous rocks, such as magnetite, ilmenite, apatite, titanite, zircon, etc., are considered by petrographers to be the first to crystallize in igneous rocks, on account of their euhedral form, and their occurrence within the silicates that make up the mass of the rock. As the effect of mineralizers is not generally recognized in developing new minerals in the igneous rocks the possibility has not been considered that these minerals form in the later magmatic stages by replacement.³⁵ Euhedral crystals of pyrite in igneous and sedimentary rocks and of garnet and magnetite in metamorphic rocks, are found unaccompanied by any adjacent alteration zone. In a similar manner we believe that euhedral magnetite, and probably other accessory minerals, are the product of mineralizers developed during the late magmatic stage.

The high-temperature minerals forming the bulk of the igneous rocks are formed at an early stage. The accessory minerals present in small amounts are not formed until a late stage and then under the influence of mineralizers. The difficulty is thus avoided as to how the magma, at an early stage, could become saturated in relatively insoluble compounds present in small amounts, and how these again could make their appearance in large amounts among the last products of consolidation.

a change might assist in clearing up misconceptions in regard to the magmatic ores and the processes of magmatic differentiation.

³⁴ Loc. cit., 102, 105.

³⁵ "If crystals of one primary mineral completely enclose crystals of another, the evidence of their relative age is absolute." (Harker, loc. cit., 179.)

STAGES RECOGNIZED IN THE FORMATION OF MAGMATIC ORES

From the above discussion it is clear that we conceive of the process of formation of plutonic rocks as consisting of stages, and that rock differentiation and ore formation are the results of an orderly series of events.

The process varies in detail with the composition and size of the individual intrusive masses undergoing crystallization. The stages in the norites and gabbros which contain the magmatic sulfid ores are as follows:

(1). The first minerals to form are olivine, the pyroxenes, and the feldspars.

(2). Magmatic alteration of the silicates often takes place prior to the formation of the ore-minerals. The most common change is that of pyroxene to hornblende (not uralite). The not uncommon hornblende gabbro, for example, may be developed by this late magmatic process, for the hornblende has probably been formed at the expense of pyroxene.

(3). Later magmatic products include interstitial pegmatitic material, interstitial quartz, and occasionally tourmaline, garnet, analcite, epidote, and calcite.

(4). The introduction of the ores by mineralizers is later, in general, than the minerals of group (3) and is unaccompanied by any secondary silicates.

(5). The pegmatite dikes, found in the neighborhood of almost all of the magmatic sulfid ores, are often later than the magmatic deposits of the basic rock itself.

(6). Hydrothermal alteration subsequent to magmatic ore deposition includes the development of chlorite, tremolite, anthophyllite, sericite, and serpentine. In general, hydrothermal alteration, altho seldom lacking, is insignificant compared with that developed in connection with deposits of other types in igneous rocks, such as those of Butte, Bingham, etc. It often does not accomplish any rearrangement of the ore, altho, in some cases, insignificant amounts of pentlandite, chalcopyrite, chalcocite and covellite are formed.

(7). At a later stage, downward enrichment and oxidation may take place. Magmatic deposits, in all cases examined, owe their metallic content to the original magmatic minerals and not to later introduced sulfids.

PART II.

DESCRIPTION OF THE VARIOUS DEPOSITS OF THE MAGMATIC ORES

GROUP I. THE NICKEL- AND COPPER-BEARING PYRRHOTITIC DEPOSITS

SUDBURY, CANADA

GEOLOGY

Those who favor the hypothesis that the Sudbury ores separated out of the magma as an early, or the earliest, constituent, and sank and collected by gravitative differentiation,⁸⁶ cite field relations as proving their contention. In spite of the voluminous literature in which these relations have been described, there are still many points in regard to which no agreement has been reached. A discussion of the genesis of these deposits is therefore incomplete without a summary of the established field relations and a statement of the problems that are unsolved as yet. Fortunately Coleman⁸⁷ has mapped in detail both the upper and lower margins of the laccolithic sheet, and given much information in regard to the structure of the individual deposits. The accompanying map (fig. 1) is drawn from a photographic reduction of his large-scale map.

As is well known, the pyrrhotite-pentlandite-chalcopyrite ore bodies of the Sudbury district occur at the base of a great sheet of "norite-micropegmatite." As shown on the map, the sheet is of unusual regularity for an intrusive mass of its size and character. It is overlain by 9000 feet of Upper Huronian sandstones, shales, conglomerates, and tuffs, which make a gentle syncline concordant in dip with the norite sheet, and underlain by a sedimentary series of great thickness (30,000 feet according to Coleman), predominantly of quartzite with included older basic irruptives. These older strata, "the Sudbury series" of early Huronian or pre-Huronian age, dip regularly to the southeast, averaging

⁸⁶ Coleman, A. P.—The nickel industry. Can. Dept. of Mines, Mines Branch. Publ. 170 (1913).

Barlow, A. E.—Geol. Surv. Can., 14, pt. H, 1-124 (1904).

⁸⁷ Loc. cit.




Fig. 1. Geologic Map of the Sudbury District. Drawn from a photographic reproduction of Coleman's latest map accompanying The Nickel Industry. Publ. No. 170, Canada Dept. of Mines.

45°. The "nickel eruptive," as the norite sheet has been named by Coleman,²⁸ therefore, has taken advantage of the unconformity between the older and younger sedimentary series, and has lifted the latter on its back without disturbing it otherwise than to develop a gentle spoon-shaped syncline.

No accurate sections across the syncline have been made, but judging from the areal map and the descriptions and sketches of the ore bodies located at the contact between the nickel eruptive and the underlying formations, the dip of the eruptive sheet at its basal contact is 40° to 45° at the surface, generally decreasing to 30° to 35° at a depth of about 1000 feet.

Coleman²⁹ has pictured in the following vivid language his concept of the manner in which the great laccolith forced its way into the sedimentary rocks:

"After the succession of sediments just mentioned had been deposited, the vast mass of molten rock of the nickel eruptive ascended, mostly from beneath an area near the middle of the southern range, as will be shown later. As the molten magma welled up from below, the crystalline rocks forming the roof of the great crucible gradually collapsed as a block twelve to fifteen miles long and several miles broad, giving rise to extensive faulting and fissuring. . . ."

"The eruptive sheet cooled extremely slowly, partly because of the great bulk of molten material and partly because of the thick mantle of sedimentary rocks above; and during the cooling much of the ore sank to the bottom, though its upper part remained mixed with norite, which finally blended into micropegmatite or granite on top, the three materials arranging themselves according to their specific gravities. . . ."

"The norite-micropegmatite sheet is one of the largest laccolithic sheets known, containing not less than 600 cubic miles at the present time, and probably having had a much greater bulk in the beginning."

Subsequent to the intrusion of the nickel eruptive, both granite masses and dikes and basic dikes were intruded along the under contact of the sill, and both "acid" and "basic" dikes cut the ore bodies.

One of the most important problems, as yet unsolved, is the origin of the "acid material," largely quartz and feldspar, frequently developing marked micropegmatitic structure, that occurs enclosed in many of the

²⁸ Coleman, A. P.—The Sudbury laccolithic sheet. *Jour. Geol.*, 15, 752-782 (1907).

²⁹ Canadian Dept. Mines, Pub. 170, p. 10 (1913).

ore bodies. Equally important is the determination of the relation of the "later granite" to the norite. Coleman has arranged the granitic rocks in three groups (see geological map, page 24). (1) Granite and granite-gneiss younger than (intrusive into) the Sudbury series and older than the norite; (2) granite mostly older than norite which was mapped without discrimination between "the older" and "the younger" groups which are of widely different origin; (3) granite younger than norite, a small area of which is shown near the Murray mine. The lack of detailed field and microscopic study prevents, for the present, the final answer to the questions raised above. Fortunately, however, Knight⁴⁰ has been studying these questions in the course of his detailed field work, and we look forward to the publication of his conclusions with great interest.

In regard to the micrographic segregations and inclusions of granitic material in the ore, we believe with Coleman⁴¹ that they are differentiates of the norite.

The micrographic structure is typical of the upper portion of the "nickel intrusive." Its occurrence in the ores as blebs and interstitial material, as well as large masses, and the variations shown in a single thin section, suggest that both the norite and the "acid" material are differentiates of a common magma.

In this regard Howe⁴² states:

"The attractive possibility has been considered that the silicious material associated with the sulphides might represent a residual portion of the magma from which the sulphides are supposed to have been derived. . . . The microscope, however, shows over and over again the absolute similarity of the acid inclusions to the foot-wall granite."

However, the foot-wall granite to which he refers may well be the "later granite." Specimens sent us by Dr. Knight from the Creighton mine, the locality which Howe studied, are decidedly micrographic. Until further data are available, the reasonable hypothesis is that the "acid" blebs, segregates and inclusions in the Sudbury ore bodies, as well as the marginal masses of the later granite, are differentiates of the parent norite magma. In many other localities, granitic differentiates are found inclosed in, or at the margins of, the norite intrusives (see the map of the Ringerike district, Norway, page 49). At Sudbury, where all the phenomena have taken place on a large scale,

⁴⁰ Loc. cit. This article contains a preliminary statement of certain of the field relations. ⁴¹ Econ. Geol., 10, 391-392 (1915). ⁴² Loc. cit., 520-521.

we should expect the "acid" differentiates to appear in large amounts. The fact brought out by Knight that dikes of the later granite cut the norite, and also that the ore rests upon a footwall of younger granite, does not prove that the ores are not magmatic. The norite and granite differentiates may be nearly contemporaneous, and the "late magmatic" ores are, in some localities, earlier than, and in other places later than, the "acid" differentiates. Evidence of extensive differentiation is found generally in rocks associated with magmatic ores.

Knight, ^{42a} has also emphasized that the Sudbury ores have wandered out into, and replaced, the schists found at the contact with the norite.

We are indebted to Dr. C. W. Knight for specimens of several types of sulfid bearing "greenstone schists." These laminated rocks consist largely of biotite, hornblende, actinolite, chlorite, clinozoisite, quartz, and plagioclase. One of these, a chlorite-actinolite gneiss from the Garson mine, is shown in fig. 10. The ore-minerals, pyrrhotite and chalcopyrite, occur in linear areas parallel to the plane of schistosity. Under the microscope the relations of the ore-minerals to the silicates are practically the same as in the igneous rocks examined. The sulfids surround the silicates, and evidently replace them to some extent, and minute sulfid veinlets definitely cut the silicates. Alteration products are practically absent.

This shows that the magmatic ores migrate to some extent into the country rocks, and apparently show the same replacement phenomena as are exhibited by the ores in the igneous rocks.

STRUCTURE OF THE ORE BODIES

Coleman, in his recent description of the Sudbury region, recognizes the following types of ore bodies:

1. *Marginal Deposits*.—These are sulfid segregations collected at the base of the main norite sheet. They gradually fade out above into barren country rock, and often show brecciation, fissuring, and "later reconcentration of ores", especially along the foot-wall, which is always a pronounced fissure. *Faulted marginal deposits* have suffered brecciation and faulting in their upper portions, and the ore has "wandered into the fissures between the blocks, either at the time as molten sulphides, or later through water transport. As chalcopyrite is everywhere the more transferable of the sulphides, it has entered the fissures more largely than the pyrrhotite. Unusually large amounts of quartz, carbonates, and

^{42a} Loc. cit.

sulphides of zinc or lead are found in these two mines (Crean Hill and Garson mines) as a result of circulating waters, and these later processes have played a larger part than in most other ore bodies of the region, whether marginal or offset."⁴³

2. *Columnar Offset Deposits*.—These remarkable ore deposits, the most notable of which occur in the Copper Cliff mine, are great cylindrical ore shoots.

"In the last report on the nickel region the Copper Cliff deposit was known to go down for 1,000 feet without interruption, as a rude oval pipe with diameters varying from 50 to 200 feet, and a dip of $77\frac{1}{2}^{\circ}$ to the northeast.

"Since that time the two ore bodies of the Victoria mine, though smaller in diameter, have been followed to the depth of 1,400 feet with no indication that they may not continue indefinitely. These two small cylinders of ore, more than 1,400 feet in length and close together, but never meeting, are not at all easy to account for on any other theory than the magmatic one, and this continuance to so great a depth was not anticipated in earlier studies of the region. . . .

"There is usually more evidence of water action than in the marginal mines, and often a certain amount of quartz and of rusty weathering carbonates is mixed with the ore, probably as later effects of magmatic waters."⁴⁴

The following ingenious explanation of these columnar ore shoots as dikes or apophyses running out from the main laccolith is favored by Coleman: ". . . it is possible that the most fluid part of the magma, the pyrrhotite-norite, entering all the fissures produced by the collapse of the underlying rock, rose from beneath under hydraulic pressure and was able, in a sense, to drill holes up through the crushed zones of rock above."⁴⁵

3. *Parallel Offsets*.—These include the great Frood-Stobie ore deposit described by Coleman as follows:

"The Frood-Stobie offset runs nearly parallel to the basic edge, but at a distance of from $\frac{3}{4}$ of a mile to $1\frac{1}{2}$ miles to the southeast. The ore more nearly resembles that of a marginal deposit than that of the ordinary offsets; and the ore body dips at an angle of 60° toward the basic edge. It is a long irregular sheet enclosing much rock, and its connexion with the edge of the norite is probably at a considerable depth below the surface. The margin of the norite parallel to it shows comparatively little ore, the sulphides belonging to it having been drained

⁴³ Loc. cit., 35.

⁴⁴ Coleman.—Loc. cit., 36-37.

⁴⁵ Loc. cit., 37.

off through a complex set of fissures to the Frood-Stobie deposit. The ore is known by diamond drilling to extend northwest beneath the country rocks to a depth of more than 1,000 feet, and at the lower points it distinctly flattens toward the basic edge of the norite. No other deposit of this type has so far been discovered; but the Frood-Stobie belt of ore is so important and so very distinct from the other types that it deserves a place by itself."

From the above descriptions it appears that the parallel offsets are mineralized dikes or sills parallel to the main foot-wall contact and dipping toward it.

From the quotations given above, it will be seen that there are evidences of hydrothermal deposition of ore and gangue minerals distinctly later than the main ore mass of the various types cited. Pyrite, marcasite, galena, sphalerite, and molybdenite are found in later veinlets, often cutting the main ore-bodies and accompanied by quartz and calcite.

MICROSCOPIC DESCRIPTIONS

We are greatly indebted to Messrs. F. L. Hess, J. F. Kemp, C. W. Knight, R. D. Longyear, F. H. Mason, M. E. Morgan, H. Ries, and T. L. Walker for specimens of rocks and ores from various mines of the Sudbury district. Our study has been facilitated by the excellent suite of Sudbury rocks obtained from the Royal Ontario Museum of Mineralogy. Our specimens are believed to be typical of the Sudbury ores, as we repeatedly find the structures and relations described by other workers.

There are three fairly distinct rock types directly associated with the ores studied by us: (1) quartz norite almost free from sulfids; (2) pyrrhotite norite with fair amounts of the sulfids and with uralitized pyroxene and some hornblende; and (3) a hornblende-bearing granitic rock with abundant sulfids constituting "the rich ore." These three types represent in a general way the three kinds of material found at the mines: (1) the "lean ore" or the barren norite, (2) the pyrrhotite norite transitional to the ore, and (3) the massive ore.

Lean Ore from the Stobie Mine.—One of the lean ores studied in detail by us is a rather fine-grained quartz norite, consisting of hypersthene, plagioclase, quartz, subordinate biotite and hornblende, magnetite, pyrrhotite, chalcopyrite, and smaller amounts of secondary chlorite and tremolite. The general relations are shown in fig. 11 (plate II).

The magnetite occurs in euhedral, subhedral, and anhedral crystals. Most petrographers would assign the magnetite to the first period of crystallization, but there is clear evidence that the anhedral magnetite

was formed later than the silicates. This is shown clearly in fig. 12. The magnetite has completely surrounded one hypersthene crystal and partly surrounded two others. All of the magnetite belongs to one generation, for there is a perfect gradation from the euhedral to anhedral forms. Thus we have evidence that the euhedral magnetite was formed during the late magmatic stage. The sulfids, pyrrhotite and chalcopyrite, which are found in occasional spots, are also later than the silicates, for with the magnetite they form hook-shaped anhedra surrounding the silicates, as shown in fig. 12. The ore-minerals occur between the silicate anhedra, and while they often surround the silicates they rarely cut across an individual crystal. Careful search, however, usually reveals a few occurrences of this sort.

Alteration products occur in the sections figured, and the natural inference would be that the sulfids are connected in some way with the alteration. Alteration of the hypersthene to tremolite and the formation of chlorite have taken place, but these minerals have been formed after the introduction of the sulfids, as figs. 13 and 14 prove. In fig. 13 a chlorite veinlet cuts pyrrhotite and chalcopyrite, and in the polished section of fig. 15 similar relations are shown. A veinlet of sulfids to the left of the chlorite veinlet of fig. 13 is accompanied by tremolite, but, as shown in figs. 27, 28, 29 and 30 (plate VI), the sulfid veinlets have no connection with alteration products. The hypersthene in the lower part of fig. 12 is partially altered, but it shows exactly the same relation to the ore-minerals as does the unaltered hypersthene in the upper part of the figure. Fig. 14 furnishes evidence that the tremolite is later than the ore-minerals. The sharp needle-crystals are not residual, but project out from the altered hypersthene into the chalcopyrite mass.

The quartz norite lean ore from the Stobie mine affords clear evidence that the ore-minerals were formed at the end of the magmatic stage, that the slight alteration of the hypersthene took place after the introduction of the ore-minerals, and that none of the sulfids have undergone rearrangement of any kind.

Pyrrhotite Norite from the Stobie Mine.—We have studied a typical specimen of pyrrhotite norite from the Stobie mine. This rock constitutes a medium grade ore with large crystals of plagioclase, aggregates of tremolite, rims of hornblende, and sulfid masses. The general relations are shown in fig. 16. The light gray areas represented in the photograph are largely aggregates of uralite needles in more or less parallel position. These aggregates are often surrounded by hornblende. The probable explanation is that hypersthene was bordered by rims of horn-

blende, and at a later stage the hypersthene, but not the hornblende, underwent uralitization. From other data we know that the sulfids were formed after hypersthene and hornblende. A study of polished sections proves that the uralitization (tremolitization) occurred after the introduction of the sulfids, and as evidence we introduce figs. 17 and 18.

Fig. 18 shows pentlandite of the first generation in veinlike areas in the pyrrhotite, and pentlandite of the second generation developing along crystallographic directions of the pyrrhotite. The rearrangement of the ores is a very minor feature in this, as in the other Sudbury specimens.

The Rich Ores.—The massive ores examined by us show residual spots of rather "acid" material, consisting largely of alkali feldspar, quartz, hornblende, and biotite. In fact the gangue of the rich ore is more like the so-called micropegmatite than the quartz norite. Microcline is abundant, and is often intergrown with quartz. In none of the rich ores have we been able to find hypersthene. Its place seems to be taken by hornblende and biotite, which are probably late magmatic alteration products. It is difficult to believe that the "acid" rock is an older foot-wall granite. It is more probably a felsic differentiate of the same magma that furnished the norite.

Rich Ore from the Stobie Mine.—Rich ore from the Stobie mine is represented by fig. 19 (polished section). The chief minerals are plagioclase, quartz, and biotite. Garnet is also present. The ore-minerals, which occur in irregular masses occasionally extending out into veinlets, replace all the silicates including garnet. The replacement along cleavage lines of biotite by the chalcopyrite is beautifully shown in thin sections. This specimen is almost entirely free from alteration products. There is, however, a little chlorite in small lath-shaped sections, and these distinctly cut the ores.

Rich Ore from the Copper Cliff Mine.—On plate I (fig. 7) we show a large polished hand-specimen from this mine. This specimen shows silicates which are believed to be residual. The residual spots contain microcline and quartz (often in graphic intergrowth), plagioclase, biotite, a little hornblende, and long acicular crystals of apatite. As far as can be told from the residual matter which has escaped replacement, the rock is a granite. The ore-minerals, magnetite and pyrrhotite, replace the silicates, especially biotite, as is shown in fig. 20. The biotite gives one the impression of being one of the last formed silicates. There are also very small biotite crystals, which may possibly belong to a second generation. The apatite is also formed at a late stage.

There are a few alteration products present, such as chlorite and sericite, but they have nothing to do with the ores.

Rich Ores from the Creighton Mine.—The Creighton is the largest mine in the Sudbury district, and fortunately our suite of specimens includes a number from that deposit. The residual silicates in the ores are plagioclase, microcline, hornblende, and biotite. Hypersthene is lacking. Quartz is abundant as an interstitial mineral, and often occurs as a quartz-feldspar intergrowth.

On account of the abundance of quartz, one of the rich ores is practically a granite or possibly a grano-diorite. Another specimen very much resembles the quartz norite in structure, but hornblende is present instead of hypersthene. It is probable that this rock was originally a norite, and this indicates the possibility not only of late magmatic minerals but also of late magmatic rocks. The same specimen that furnished the thin section showing the norite texture also shows microcline and a subgraphic intergrowth of quartz and feldspar. This variation in the constituents within a small space is a characteristic of the igneous rocks containing magmatic ores.

A photograph of rich ore from the Creighton mine is shown in fig. 8 (plate I). This specimen furnished us the photomicrographs of figs. 25, 27-30, 32, and 33. A similar specimen furnished photomicrographs 23, 24, 31, and 34.

Polished sections showing the general relation of the sulfids to the silicates in these specimens are shown in figs. 27, 28, 31, and 32. Thin sections, which are necessary for the identification of the silicates, are represented by figs. 23 and 25.

The sulfids are later than the silicates. This is almost certain from the irregular hook-shaped anhedral which surround and occasionally project into the silicates; but if any doubt exists as to the later origin of the sulfids note the veinlets in plates V, VI, and VII. The veinlets replace almost indiscriminately the various silicates (see figs. 24, 28, and 30), but certain sulfid areas show marked selective replacement of the feldspar of the quartz-feldspar intergrowth, as in fig. 31. For an especially good illustration of replacement veinlet see fig. 24, where fresh biotite has been invaded by the sulfid. In fig. 26 a hornblende crystal has been cut squarely in two by a sulfid veinlet. Note the branching veinlets of figs. 28 and 30. The only explanation of these veinlets is that they are later than the silicates. The proponents of the magmatic theory for the Sudbury ores have ascribed these veinlets to rearrangement, and thus they reconcile their theories with microscopic work such

as that of Dickson. That the sulfid veinlets are of the same generation as the main sulfid masses and are not due to later rearrangement is shown in fig. 29, where the sulfids extend out into a veinlet without any break in the continuity or other evidence of later origin.

The late origin of the sulfid veinlets can hardly be doubted, but some may be inclined to suggest a hydrothermal origin for them. This is disproved by our findings. For example in the thin sections of the Creighton ore we found tremolite (and possibly talc) pseudomorphous after original hypersthene. The pseudomorphs contain minute magnetite crystals which were formed by alteration. The pseudomorphs are cut by veinlets of chalcopyrite, but the chalcopyrite has not wandered into the cracks of the tremolite. This shows that the hydrothermal alteration of hypersthene to tremolite (or talc) and magnetite occurred after the introduction of the chalcopyrite.

That the magnetite, as well as the sulfids, is formed at a late magmatic stage is indicated by fig. 20. The magnetite contains included ilmenite plates as illustrated by fig. 21. This magnetite-ilmenite intergrowth is a characteristic feature of magmatic ores.

Polished sections of massive, almost solid, ore from the Creighton mine are represented by figs. 22, 35, and 36. Fig. 22 shows an area of silicates which is doubtless a relict of the same granitic material present in the other Creighton samples. Magnetite has been replaced by the sulfids and the pentlandite has replaced pyrrhotite in vein-like masses. Veins of pentlandite are well shown in fig. 35; also minute tufts of brush-like crystals of a pale-yellow mineral, probably pentlandite of a second generation. Fig. 36 is a highly-magnified view of one of these crystals which extends out from a veinlet of chalcopyrite, probably of the second generation. Figs. 35 and 18 show what a minor amount of rearrangement has taken place in the typical magmatic ore of Sudbury.

At some of the Sudbury mines, notably the Worthington, certain sulfids which are not typically magmatic have been found. Among these minerals are pyrite and polydymite, but many others have been reported.⁴⁶

We have examined several Sudbury ores containing pyrite. In fig. 37 is represented a supposed specimen of polydymite from the Vermilion mine. This contains pyrite in the form of veinlets evidently formed at a late stage. The relation of the polydymite to the magmatic sulfids is not entirely certain, but the examination of a very fresh speci-

⁴⁶ Barlow, A. E.—The nickel and copper deposits of Sudbury, Ontario. Geol. Surv. Can. Ann. Rept. 14, pt. H, 93 et seq.

- The Sudbury area. 12th Inter. Geol. Cong. Guide-Book no. 7, 48 pp. (1913).
- A classification of the Sudbury deposits. *Trans. Can. Min. Inst.*, 16, 283-288 (1913).
- Discussion of (Petrographical notes on the Sudbury nickel deposits—Howe). *Econ. Geol.*, 10, 390-393 (1915).
- Dickson, C. W.—Note on the condition of nickel in nickeliferous pyrrhotite from Sudbury. *Eng. and Min. Jour.*, 73, 660 (1902).
- Note on the condition of platinum in the nickel-copper ores from Sudbury. *Am. Jour. Sci.*, (4) 15, 137-139 (1903).
- The ore deposits of Sudbury, Ontario. *Trans. Am. Inst. Min. Eng.*, 34, 3-67 (1909).
- Garnier, J.—Mines de Nickel, Cuivre, et Platine du District de Sudbury, Canada. *Mem. Soc. des Ing. Civiles*, Paris, 1891.
- Goodwin, W. L.—*Am. Jour. Sci.*, (3) 47, 312-314 (1894).
- Gregory, J. W.—Origin of the Sudbury nickel ores. *Geol. Mag.*, (new series) 5, 139-140 (1908).
- Hixon, H. W.—Geology of the Sudbury district. *Eng. and Min. Jour.*, 79, 334-335 (1905).
- The ore deposits and geology of the Sudbury district. *Jour. Can. Min. Inst.*, 9, 223-235 (1906).
- The Sudbury nickel region. *Eng. and Min. Jour.*, 82, 313-314 (1906).
- Hore, R. E.—Origin of the Sudbury nickel and copper deposits. *Min. and Eng. World*, 36, 1345-1349 (1912).
- Magmatic origin of Sudbury nickel-copper deposits. *Mich. Geol. and Biol. Surv.*, Pub. 16, pp. 11-37 (1914).
- Howe, E.—Petrographical notes on the Sudbury nickel deposits. *Econ. Geol.*, 9, 505-522 (1914).
- Kemp, J. F.—An outline of views held today on the origin of ores. *Mineral Industry*, 4, 755 (1895).
- Knight, C. W.—Origin of Sudbury nickel-copper deposits. *Eng. and Min. Jour.*, 101, 811-812 (1916).
- Miller, W. G.—On some nickeliferous magnetites. *Brit. Assn. Adv. Sci.*, Rept. 1897, pp. 660-661 (1898).
- Miller, W. G., and Knight, C. W.—Sudbury, Cobalt, and Porcupine geology. *Eng. and Min. Jour.*, 95, 1129-1133 (1913).
- Penfield, S. L.—On pentlandite from Sudbury, Ontario, Canada, with remarks upon three supposed new species from the same region. *Am. Jour. Sci.*, (3) 45, 493-494 (1893).
- Silver, L. P.—The sulphide ore bodies of the Sudbury region. *Jour. Can. Min. Inst.*, 5, 528-551 (1902).
- St. Clair, S.—Origin of the Sudbury ore deposits. *Min. and Sci. Press*, 109, 243-246 (1914).
- Stewart, L.—The Creighton mine of the Canadian Copper Co., Sudbury Dist., Ontario. *Jour. Can. Min. Inst.*, 11, 567-585 (1908).
- Stokes, R.—The Sudbury nickel-copper field. *Min. World*, 27, 637-639, 799-801 (1907).
- Stutzer, O.—Die Nickelerzlagertstätten bei Sudbury im Kanada. *Zeit. f. prakt. Geol.*, 16, 285-287 (1908).

- Thomas, K.—The Sudbury nickel district, Ontario, Canada. *Min. and Sci. Press*, 105, 433 (1912).
- The Sudbury nickel district of Ontario. *Eng. and Min. Jour.*, 97, 152-154 (1914).
- Thompson, P.—The Sudbury nickel region. *Eng. and Min. Jour.*, 82, 3-4 (1906).
- von Foullon, H. B.—Über einige Nickelerzvorkommen. *Jahr. d. k.k. geol. Reichsanstalt*, 42, 223-310. Vienna (1892).
- Walker, T. L.—Notes on nickeliferous pyrite from the Murray mine, Sudbury, Ont. *Am. Jour. Sci.*, (3) 47, 312-314 (1899).
- Geological and petrographical studies of the Sudbury nickel district. *Quar. Jour. Geol. Soc.*, 53, 40-66 (1897).
- Certain mineral occurrences in the Worthington mine, Sudbury, Ontario, and their significance. *Econ. Geol.*, 10, 536-543 (1915).
- Williams, G. H.—Notes on the microscopical character of rocks from the Sudbury mining district. *Can. Geol. Surv. Reports*, (new series) 5, pt. I, Report F (1893).

THE ALEXO MINE, ONTARIO, CANADA

GEOLOGY

This occurrence of nickeliferous pyrrhotite, twenty miles southeast of Porcupine, Ontario, has been mentioned briefly by Coleman,⁸⁰ and described in detail by Uglow.⁸¹ The geological relations are not disclosed on account of lack of exposure, and the mineralogical relations are complicated by an intense serpentinization. The ore occurs as a lens in serpentinized rock at the contact with rhyolite. In cases of this kind microscopic work is especially valuable. The unaltered rock is considered by Uglow to have been a peridotite of the wehrlite or harzburgite variety. The only exposure is in the immediate vicinity of the ore body, and developments are not sufficient to show the structural relations.

UGLOW'S CONCLUSIONS

Uglow examined the ore in both thin and polished sections, has shown the relations in photographs, and has brought out in a convincing way the phenomena of replacement as shown by his microscopic study. He states that the ore (pyrrhotite, pentlandite, chalcopyrite) "eats its way through the matrix of the serpentine" . . . "replaces it, forming a network of ore" . . . "extends between the crystals into fractures and cracks in the latter" . . . that "ore replaces part or all of an olivine crystal" and produces pseudomorphs—"magnetite pseudomorphs

⁸⁰ Coleman, A. P.—The Alexo nickel deposit. *Jour. Geol.*, 5, 373-376 (1910).

⁸¹ Uglow, W. L.—A new nickel occurrence in northern Ontario. *Jour. Can. Min. Inst.*, 14, 657-677 (1911).

which have resulted from the alteration of the olivine, have become partly or wholly replaced by pyrrhotite," etc.

He finds that the order of the formation of the ore-minerals is (1) magnetite, (2) pyrrhotite, (3) pentlandite and chalcopyrite; and remarks: "It is difficult to conceive of the sulphides as differentiations from a molten magma when they are as a matter of fact deposited one after the other, the younger ones occurring as vein-like masses in the older."

His work is incomplete in that he does not investigate what portion of the phenomena of replacement and ore migration is to be connected with serpentinization, and what portion with the first deposition of the ores prior to serpentinization. Lindgren⁵² states: "Here, as in so many other cases, secondary changes seem to have been confused with primary deposition."

We have, therefore, investigated in detail the relative amount of transfer and migration that is connected with serpentinization, and have been able to show that the phenomena of replacement mentioned above antedate serpentinization, and are connected with the original deposition of the ores.

MICROSCOPIC DESCRIPTION

Polished sections of ore of the Alexo mine kindly furnished by Mr. F. H. Mason of the Canadian Government Exhibition Commission, show pyrrhotite with minor amounts of pentlandite and chalcopyrite in a gangue of serpentine. The serpentine is the result of alteration of olivine. The general relations of the minerals are shown in fig. 39. Residual cores, apparently of olivine, prove to be serpentine upon examination of the thin sections. See also fig. 40, which is an enlargement of a portion of fig. 39. The section shown in figs. 39 and 40 was polished to bring out the silicates. On the other hand, the surface shown in figs. 41-44 was polished especially for the sulfids. Pentlandite occurs in fair amounts rather evenly distributed through the ore, with a tendency to alignment along crystallographic directions of the pyrrhotite which it replaces. The other sulfid present is chalcopyrite, which occurs in two, or possibly three, generations. In fig. 41 there is a little magmatic chalcopyrite and pentlandite. The serpentinization has been accompanied by a migration of nickel to form a second generation of pentlandite along veinlets (fig. 42), and of copper to form a second generation of chalcopyrite (fig. 44). The minute euhedral crystals of chalcopyrite shown in fig. 44 may pos-

⁵² Lindgren, W.—Mineral Deposits, 765.

sibly represent a third generation of chalcopyrite. Altho there has been extensive alteration of the silicates in the Alexo ore, microscopic study shows clearly (see especially fig. 43) that this alteration was subsequent to the main period of ore formation and that the migration of ore during serpentinization was relatively slight.

SUMMARY

Summarizing the stages in the Alexo ore deposit we have (1) the formation of pyrrhotite, pentlandite, and chalcopyrite in the order named, at the end of the magmatic period, probably by selective replacement of the ground-mass of a picrite (pseudo-porphyrific peridotite); (2) the alteration of olivine and possibly of other silicates to serpentine; (3) accompanying the serpentinization there was a slight migration of copper and nickel to form second generations of chalcopyrite and pentlandite respectively, and also the formation of magnetite in very minute crystals and veinlets.

BIBLIOGRAPHY OF THE ALEXO ORE DEPOSITS

- Coleman, A. P.—The Alexo nickel deposit. *Econ. Geol.*, 5, 373-376 (1910).
 —The nickel industry with special reference to the Sudbury region, Ontario. Can. Dept. Mines, Mines Branch. Report 170, pp. 112-114 (1912).
 Uglow, W. L.—The Alexo nickel deposit, Ontario. *Ont. Bur. Mines. Ann. Rept.* 20, pt. 2, 34-39 (1911).
 —The Alexo mine. A new nickel occurrence in northern Canada. *Jour. Can. Min. Inst.*, 14, 657-677 (1911).
 —Summary report on the Sudbury nickel field. Can. Dept. Mines, Mines Branch, Summary Rept. 1911, pp. 87-89.

THE FRIDAY MINE, SAN DIEGO COUNTY, CALIFORNIA

The occurrence of nickel ores in the Friday mine, San Diego county, California, was reported by Merrill^{52a} and described by Calkins,^{52b} whose paper furnished the geological data summarized in the following paragraphs:

The mine is four miles from the town of Julian, which lies half-way between San Diego and the Salton Sea. It is located on the crest of the broad range which forms the northern extension of the Cordillera of Lower California.

^{52a} Merrill, F. J. H.—Geology and Mineral Resources of San Diego and Imperial counties. California State Mining Bureau Report, biennial period 1913-1914, p. 40.

^{52b} Calkins, F. C.—An occurrence of nickel ore in San Diego county, California. *S. Bull.* 640, U. S. Geol. Surv., 77-82 (1916).

The complex batholith of Lower California extends into San Diego county, and the granitic rocks of the region are probably closely related to the larger masses to the south. Metamorphosed sediments are included within the granite, and near the mine a mass of gabbro of unknown size lies in contact with mica schist.

The ore occurs as a shoot or lens at the contact of the gabbro with the schist. The latter dips steeply southward, and the ore lies at the base of the gabbro. Considerable fracturing is reported in the vicinity of the ore body, and the ore is penetrated by a small pegmatite dike containing conspicuous crystals of tourmaline.

MICROSCOPIC DESCRIPTION

Thru the kindness of Mr. Beecher Sterne, president of the Friday Copper Mines Company, we obtained a suite of rocks and ores from the Friday mine, including specimens from the recently developed lower levels. The rock is an olivine gabbro with plagioclase (Ab_8, An_7 in one section), olivine, and both orthorhombic and monoclinic pyroxene. A pale brown hornblende occurs as rims around the border of, and as patches within, the pyroxene. The hornblende is in parallel position with the pyroxene, and is doubtless a magmatic alteration product. Small tremolite prisms replace the pyroxene and plagioclase. Another alteration product is calcite, which occurs in veinlets and occasionally in zonal crystals.

The polished sections of the massive ores consist largely of pyrrhotite with residual spots of silicates. Associated with the pyrrhotite is a considerable amount of pentlandite. Calkins reported this as polydymite, but it has the characteristic cleavage, relief, and color of pentlandite. Chalcopyrite in small amounts is also present in the sections, and is distinctly later than the pyrrhotite and pentlandite. A second generation of pentlandite is developed along cracks in the pyrrhotite. A brass-yellow mineral occurs in veinlets and reticulate masses as a replacement of pyrrhotite. This was called pyrite by Calkins, but more probably it is marcasite. The marcasite gives the impression of being a very late mineral. It is usually extensively developed along calcite veinlets which occur as a net of intersecting stringers cutting all the other minerals. The polished sections also show that tremolite is later than the sulfids.

SUMMARY

In the Friday deposit we have the following sequence of events: (1) The crystallization of olivine, pyroxenes, and plagioclase. (2) A slight development of hornblende by magmatic alteration. (3) The

formation of pyrrhotite, pentlandite, and chalcocopyrite in the order named by the replacement of the above mentioned silicates. (4) The development of tremolite as a hydrothermal mineral and the development of pentlandite of a second generation in cracks in pyrrhotite. (5) The extensive development of calcite and marcasite veinlets.

With the exception of the marcasite and calcite this deposit is similar in mineral composition and paragenesis to magmatic deposits of this type.

THE GOLDEN CURRY MINE, ELKHORN, MONTANA⁵³

GEOLOGY

An interesting deposit of magmatic gold- and copper-bearing pyrrhotite (non-nickeliferous, however) occurs in a marginal "basic" segregation of the Boulder quartz monzonite batholith. The deposit thus differs from the normal type of magmatic pyrrhotite deposits in respect to the geological relations and the character of the mother rock. The mineral composition, the inclusion of the ore in a subsilicic rock, and the relatively unaltered condition of the ore and country rock, give this deposit the characteristics of a magmatic ore. It is possibly the unmetamorphosed equivalent of the lenses of pyrrhotite in "basic" layers of gneiss, and is an indication that certain of these puzzling occurrences are altered magmatic segregations.

The Boulder batholith is well known to students of ore deposits because it was accompanied and followed by extensive mineralization, and especially because it encloses the enormous copper deposits at Butte.

The Golden Curry mine at Elkhorn is in a region of intense mineralization,⁵⁴ which occurred in and near the roof of the batholith. In this property there are two types of ore deposits: (1) a contact deposit between quartz monzonite and limestone, consisting of magnetite and some chalcocopyrite accompanied by garnet; (2) the magmatic pyrrhotite deposit. The latter occurs 250 feet from the contact mentioned above, as the sulfid rich portion of a lens of fresh monoclinic pyroxene, pyrrhotite, and chalcocopyrite. This is surrounded by a border zone, which grades into the normal quartz monzonite by the addition of, first, plagioclase, then hornblende, quartz, and the accessory minerals of the quartz monzonite, and finally biotite and orthoclase. According to

⁵³ Knopf, A.—A magmatic sulphide ore-body at Elkhorn, Mont. *Econ. Geol.* 8, 323-336 (1913).

⁵⁴ Knopf, A.—Ore deposits of the Helena mining region, Montana. *Bull.* 527, U. S. Geol. Surv. (1913).

Knopf, the deposit is characterized by the absence of pneumatolytic or hydrothermal alteration products. In the vicinity, however, intense pneumatolytic action is shown in the Queen mine, the ore of which is argentiferous galena, accompanied by quartz and tourmaline. Strong mineralization of a later cooler phase is also represented in the highly productive Elkhorn mine. The ore is a replacement deposit in dolomite, consisting of argentiferous galena unaccompanied by metasomatic gangue minerals.

The Boulder batholith develops in places a border zone distinctly more basic than the normal quartz monzonite. The ore deposits, however, with the exception of the Golden Curry mine, are connected with the "acid" differentiation products, as emphasized by Billingsley.⁵⁵ Aplite masses, aplite and pegmatite dikes, often accompanied by tourmaline and sulfids, are well developed in and near the roof of the batholith. Contact deposits are common, with axinite, tourmaline, etc., in addition to the usual minerals. Tourmaline-copper and tourmaline-lead-silver ores are the high-temperature phases brought about by the mineralizing action of the intrusive. It is therefore of considerable interest to know whether the "basic" segregation of the Golden Curry mine was formed at an early or a late stage. Was there a gap between the magmatic deposits and the later high-temperature phase so extensively developed in this region?

MICROSCOPIC DESCRIPTION

Mr. Adolph Knopf of the United States Geological Survey has kindly furnished us with thin sections of the pyrrhotite-augite ore from the Golden Curry mine. The rock is practically a pyroxenite, consisting mainly of monoclinic pyroxene (augite) and pyrrhotite, with minor amounts of hornblende, tremolite, and chalcopyrite.

The replacement of pyroxene by the pyrrhotite and chalcopyrite is well shown in fig. 45. The ore-minerals not only surround the pyroxene, but occasionally cut straight across a pyroxene crystal, as shown a little to the left and below the center of the photograph. The replacement of the pyroxene by the sulfids is also proved by fig. 47, for the small pyroxene crystal near the center is cut in two by a veinlet. The pyroxene has been altered to green hornblende (not uralite) in occasional patches (the darker gray spots in figs. 45 and 47). Definite proof that the sulfids are later than the hornblende and replace it along cleavage lines is shown in the lower portion of fig. 46. Tremolite is

⁵⁵ Billingsley, P.—The Boulder batholith of Montana. *Tras. Am. Inst. Min. Eng.*, 51, 31-56 (1915).

formed at a still later stage, for as shown in fig. 47, it projects from the end of a hornblende crystal into the sulfids.

We agree with Knopf that the pyrrhotite-chalcopyrite ore of the Golden Curry mine is a magmatic deposit showing very little post-mineral alteration, but we believe that the sulfids were formed at a late magmatic stage, and after the partial magmatic alteration of pyroxene to hornblende.

PROSPECT HILL, LITCHFIELD, CONNECTICUT

An interesting series of sulfid-bearing igneous rocks varying from gabbro and norite to peridotite and pyroxenite was described by Hobbs⁵⁶ from this locality. Extreme magmatic differentiation has produced a great variety of rock types within a small area.

Recently Howe⁵⁷ has studied the relation of the sulfids to the silicates in these rocks. He identifies pyrrhotite, pentlandite, and chalcopyrite, which were formed in the order named. A prominent feature is the presence of hornblende and biotite as magmatic alteration products of pyroxene, but no mention is made of any post-magmatic alteration. Howe concludes that the ore-minerals mentioned are magmatic sulfids which separated for the most part at an early stage in the cooling of the magma, but remained liquid until the silicates had crystallized. He believes, however, that the sulfid veinlets in the silicates were formed by the replacement of the latter.

Dr. Howe has kindly sent us some specimens of these interesting sulfid-bearing rocks, and we have corroborated nearly all of his findings. The Litchfield sulfids are magmatic sulfids, if such exist. There is no evidence of any rearrangement of the ore-minerals. The veinlets do not belong to a later generation, and all the ore-minerals are late magmatic. Altho the rocks are exceptionally fresh, perhaps more so than the average unmineralized igneous rock, we find some serpentinization of the olivine. This is especially well shown in the polished sections.⁵⁸

We find that there has been slight post-magmatic alteration of the silicates, but conclude with Howe⁵⁹ "that the relations of the sulphides to one another were . . . determined during the magmatic period."

⁵⁶ Festschrift Harry Rosenbusch, 25-48. Stuttgart, 1906.

⁵⁷ Howe, E.—Sulphide-bearing rocks from Litchfield, Conn. *Econ. Geol.*, 10, 330-347 (1915).

⁵⁸ The alteration products are very apparent in well polished sections, tho of course the identity of the minerals must be determined in thin sections or crushed fragments.

⁵⁹ Loc. cit., 339.

KNOX COUNTY, MAINE

Bastin⁶⁰ has described in detail a pyrrhotitic peridotite from this locality. The peridotite is thought to be a differentiate of granite. Chalcopyrite and pyrite are associated with the pyrrhotite, but the relation of the pyrite to the other ore-minerals is not discussed. Hornblende is locally important, and occurs both as crystals and as rims between feldspar and olivine. The secondary alteration (serpentinization) is later than the ore. His photographs show pyrrhotite surrounding the olivine and penetrating it in embayments.

This is evidently a typical magmatic deposit, with the ore later than the primary silicates.

MOUNTAIN, WISCONSIN⁶¹

This deposit occurs in a basic dike 60 to 200 feet wide. The ore is massive pyrrhotite which "merges into gabbro on indefinite lines." The author does not give the results of a careful field study, and no microscopic investigation of the rock or ore is recorded.

OTHER PYRRHOTITE DEPOSITS IN THE UNITED STATES

In addition to its occurrence in the magmatic ores, pyrrhotite is also found in other high-temperature deposits. Among these are the lenticular ore-bodies found in gneisses and schists. Those in the basic portion of the gneiss may be magmatic in origin, and those in the acid layers may be related to "acid" or pegmatitic extracts. Altho some of the pyrrhotite deposits occurring in metamorphic rocks, for example that of the Gap mine, Lancaster, Pennsylvania,⁶² have been assigned to the magmatic group, most of them have been so modified by intense metamorphism that they no longer show the characteristics of typical magmatic deposits.

INSIZWA RANGE, EAST GRIQUALAND, SOUTH AFRICA

Our knowledge of the deposits near the town of Mount Ayliff, in Cape Colony, is due chiefly to the excellent, tho brief, report of Du

✓ ⁶⁰ Bastin, E. S.—A pyrrhotitic peridotite from Knox county, Maine. Jour. Geol., 16, 124-138 (1908).

✓ ⁶¹ Bagg, R. M.—The discovery of pyrrhotite, with a discussion of its probable origin by magmatic differentiation. Econ. Geol., 8, 369-373 (1913).

✓ ⁶² Kemp, J. F.—The nickel mine at Lancaster Gap, Pennsylvania, and the pyrrhotite deposits at Anthony's Nose, on the Hudson. Trans. Am. Inst. Min. Eng., 24, 620-633 (1894).

Toit.⁶³ He has discussed in a satisfactory manner the general geology and the petrography of the region. His microscopic description of the ores is brief, and it is chiefly in this respect that further information is needed. Unfortunately we have not been able to procure specimens from these deposits.

The deposits show certain similarities to those of the Sudbury district. The ore-minerals are the same, and they occur at the base of a large "basic" sill. They differ from the Sudbury deposits in certain respects, however, and features that are hidden at Sudbury are exposed by the erosion of the mountain mass of the Insizwa range. Du Toit's results are summarized in the following paragraphs.

Interest has been aroused in these deposits, first, by the discovery of copper, then nickel, and recently of platinum. The ores consist of pyrrhotite, pentlandite, chalcopyrite, with some pyrite, niccolite, and a platinum mineral, probably sperrylite. The ores occur at the base of the largest of a series of sills of olivine gabbro and olivine norite. The country rock, therefore, is more "basic" than the Sudbury "nickel intrusive." At this locality an extensive series or group of sills occurs in the Beaufort shales and sandstones of the Karoo system. The sills are flat-topped, but the under surface is funnel-shaped, narrowing down into the feeding dikes. Strange to say, the intruded shales are not tilted but are "undisturbed and almost absolutely flat, bounded by an under regularly curved surface of fractures."

The ores are intimately mixed with the silicates and "intrude" the country rock to some extent. The silicates of the sill show no alteration where they adjoin the ores. In addition to the ores, a granitic extract has been segregated along the base of the sill. This occurs as a network of dikes and dikelets, from a fraction of an inch to a foot in thickness. Intense contact action has affected the sedimentary rocks, and seems to be closely related to the dikes. The product of the contact metamorphism is a hornfels, described as a quartz-cordierite-feldspar rock with abundant biotite and enstatite in places. In the calcareous layers epidote, zoisite, diopside, enstatite, wollastonite, and garnet are developed.

Du Toit's figure⁶⁴ shows relations similar to those we have found in the magmatic ores from other localities, and which we interpret as indicating a replacement of the gangue by the ore-minerals. Du Toit

⁶³ Du Toit, A. L.—Report on the copper-nickel deposits of the Insizwa, Mount Ayliff, East Griqualand, Cape of Good Hope. Dept. Mines, 15th Ann. Rept. of the Geol. Comm., 111-142 (1910).

⁶⁴ Loc. cit., fig. 4, p. 139.

makes the order of formation of the sulfids the reverse of that determined by Campbell and Knight for the Sudbury deposits. An examination of Du Toit's sketch⁶⁵ shows the same marginal relations of chalcopyrite to the other sulfids as found in Sudbury, and we venture to suggest that further microscopic study of the ore will show that the order of the formation of the sulfids is the same at Mount Ayliff as in all other known deposits of this type.

Summarizing, we find the following suite of events connected with intrusion and ore formation at Mount Ayliff:

(1) The intrusion of a complex family of "basic" sills connected by cross-cutting dikes.

(2) The crystallization of the silicate minerals of the gabbro and norite.

(3) The development of the sulfid ores in certain localities in the basal portion of the largest of the sills by replacement of the silicates, but unaccompanied by the development of secondary silicates.

(4) The squeezing out of a granitic extract, forming a complex set of small dikes at the base of the sill, and the development of intense contact metamorphism due to gases accompanying the "acid extract." Ore formation also occurred in this stage, as ore-minerals are found in the pegmatite and granitic dikes.

NORWAY

GEOLOGY

In Norway, Nature has constructed a series of small-scale models to illustrate the type of nickel, copper, and iron sulfid deposits of magmatic origin. These instructive examples have been described in detail by Vogt,⁶⁶ whose classic work has established the existence and characteristics of this type of ore deposits. In as much as a part of the microscopic data admit of explanation according to Vogt's hypothesis of an intrusive sulfid magma, we discuss in some detail his conclusions as to the origin of these ores.

In many of the Norwegian occurrences neither the ore nor the norite has suffered metamorphism, and the geological relations are simple, so that it seems possible to select the phenomena that are characteristic of this type of deposits. It is essential for an understanding of the problems that this be done, for in many of the deposits of nickeliferous pyrrhotite elsewhere the data are either wanting or not easily

⁶⁵ Loc. cit., fig. 3, p. 137.

⁶⁶ Cited, p. 5.

deciphered. At Sudbury, for instance, it is difficult to compile the story of ore formation from the geologic records, overcrowded with the details of the eventful igneous and metamorphic history of which ore deposition is a mere episode. There, inclusions of mafic material may be explained either as "basic" segregations or as inclusions of "the older norite," as best suits the hypothesis favored by the individual investigator. Felsic masses and dikes are variously explained as inclusions of an older granite, as "acid" segregations of the norite, or as a distinctly younger intrusive granite.

In addition to the little altered types, some of the Norwegian deposits are modified by subsequent metamorphism, and in others the ores are accompanied by metasomatic borders of garnet (fig. 5). Hydrothermal alteration with the formation of pyrite appears to be developed where pegmatitic intrusions are numerous and active mineralizers were present (fig. 3).

These gradation types are fully as instructive as the normal occurrences, and support our contention that the typical magmatic deposits are one phase of high-temperature deposition, and changes in the character of the mineralizers result in gradations towards the other types.

In Norway there are about fifty deposits of pyrrhotitic copper-nickel ores which occur exclusively at or near the margins of small norite or gabbro stocks. The stocks are round or elliptical in outline, and some are funnel-shaped in cross-section. Figs. 2 and 3, taken from Vogt, illustrate typical occurrences. The individual intrusives are small, rarely exceeding a few hundred feet in diameter, and the amount of ore in each stock is roughly proportional to its size.




Fig. 2. Meinkjær mine, Norway. (Distance across sketch about 500 ft.) (After Vogt, *Zeit. f. prakt. Geol.*, plate VI, fig. 3, opposite p. 133, Jahrgang 1893.)

PETROGRAPHY AND MINERALOGY

The ore-bearing rock may be classed broadly as norite (\pm quartz, olivine, and diallage), and in a few localities it is so extensively uralitized as to be classified by some authorities as a "gabbro-diorite." Each individual stock has undergone extensive differentiation, especially in the vicinity of the ore bodies. The basic segregations are mixtures of bronzite, hornblende, olivine, and biotite. They occur in masses, smaller inclusions, and, at Romsaas, as a peculiar orbicular norite. The felsic




Fig. 3. Erteli mine, Ringerike, Norway. (After Vogt, *Zeit. f. prakt. Geol.*, fig. 8, plate V, Jahrgang 1893.)

products of differentiation are as prominent as the mafic, and occur as masses of granite, streaks and dikes of pegmatitic and aplitic material, and small marginal veinlets of aplite and pegmatite. A similar differentiation appears to be a characteristic of all deposits of this class. The margins of the stocks often show severe fracturing and brecciation, and the ore may cement these igneous breccias, as at Sudbury.

According to our ideas, it would appear that (1) the "basic" portions of the norite solidified by the sinking of crystals; (2) portions of this "basic" material were caught up as inclusions in the normal norite, especially along the margins of the stocks; (3) the main mass of the rock crystallized as norite; (4) felsic dikes and masses were squeezed out as marginal segregates during a period of peripheral brecciation, which facilitated the concentration and escape of mineralizers from the deeper portions of the magma; (5) ore deposition, at a late stage, was preceded, accompanied, and followed, by the "acid" secretions, as the ore is found in the felsic dikes, is later than the silicates of the same, and the main ore masses are cut by the dikes.




Fig. 4. Detailed cross-section of ore-body at the Meinkjær mine, Norway. (After Vogt, *Zeit. f. prakt. Geol.*, fig. 29, p. 136, Jahrgang 1893.)

The ores occur chiefly at the margins of the stocks, and to a minor amount as segregations within the norite, and also as impregnations in the schists and gneisses in which the norite masses are enclosed. As has been mentioned, the ores are further localized where marked differen-



Fig. 5. Pyrrhotite gabbro from Erteli mine, Norway. Pyrrhotite veinlet cutting feldspar and diorite with rim of garnet. (x 100.) (After Vogt, *loc. cit.*, p. 139.)



Fig. 6. Pyrrhotite veinlets cutting hornblende. Flaad mine, Evje, Norway. (x 50.) (After Vogt, *loc. cit.*, p. 139.)

tiation has taken place. At the Flaad mine, especially, the ores are closely related to felsic differentiates.

The ores show the same group of minerals (chiefly magnetite, pyrrhotite, pentlandite, and chalcopyrite), and the same relation to each other and to the schists, as we have found in the deposits studied elsewhere. Pyrite is mentioned by Vogt as of common occurrence in these

deposits. He describes it especially in the Erteli mine at Ringerike, and the Flaad mine at Evje. Ores from both these localities, furnished through the courtesy of Professor J. F. Kemp, were examined by us. The pyrite from Ringerike occurs as euhedral and irregular grains, cutting both the other ore-minerals and the silicates. Its position in the sequence of mineral formation could not be definitely determined from the material available. Pyrite, in the ore from Evje, however, clearly cuts garnet and also all the ore-minerals, including chalcopyrite, as groups of anastomosing sub-parallel veinlets (fig. 48). These veinlets widen into irregular areas, and even subhedral crystals are often developed. This makes it probable that the euhedral and subhedral pyrite crystals of fig. 48, as well as the veinlets, are later than the pyrrhotite. Chlorite and calcite cut the sulfids in sharp veinlets, and occur along the pyrite veinlets. They are certainly later than all the sulfid minerals except pyrite, and may be later than it.

As already stated, pyrite is not commonly found in the magmatic ores. At Sudbury, it is found especially in the Worthington offset, connected with later veins of hydrothermal origin. In the Ringerike deposits ore has been mined from disseminations in the country rock, and the pyrite in these ores may have been developed in the schists prior to the introduction of the ores. At Evje, as stated above, the pyrite is the last sulfid mineral to form, and may have been developed during the extensive uralitization of the country rock. A complete microscopic study of all the Norwegian ores according to modern petrographic and metallographic methods is needed in order to determine the paragenesis of the pyrite.

As described and figured by Vogt (fig. 5), veinlets of sulfid ores are accompanied by garnet, showing a gradation of these deposits to other high-temperature deposits.

Contrary to the general impression, Vogt states that the ores in all cases are later than the silicates. He notes⁶⁷ the frequent occurrence of microscopic veinlets in the cracks and cleavage planes of the silicate minerals. We wish to emphasize again, that these data, checked by us, admit of only two explanations, viz., (1) that of Vogt, who postulates an intrusion of molten sulfids into the solidified rock; or (2), our hypothesis that the sulfids were formed by the replacement of silicates under the influence of mineralizers. This process, however, belongs to a late magmatic period, and not to destructive pneumatolytic or hydrothermal stages, and the deposits, therefore, may properly be designated as magmatic.

⁶⁷ Zeit. f. prakt. Geol., Jahrgang 1893, p. 139.

DISCUSSION OF VOGT'S CONCLUSIONS

In the light of the data summarized in the preceding pages we may now examine in detail the data listed by Vogt,⁶⁸ in his latest contribution, as favoring his hypothesis.

"As suggested more particularly by Vogt in 1893, the nickel-pyrrhotite deposits are to be regarded as magmatic segregations in gabbro or in chemically analogous dike rocks. The following points speak for such an origin:

1. The connection of the numerous deposits in different countries with occurrences of gabbro or exceptionally with equivalent dike rocks, is constant and regular.

2. The several occurrences so resemble one another not only geologically but also mineralogically that they must be of the same genesis throughout, while the unvarying character of the deposits postulates a simple process of formation.

3. Gradations between ore and gabbro through the intermediate stage of pyrrhotite-gabbro are often to be observed, and therefore the ore essentially must have been formed in a similar manner to that rock.

4. The structure of the clean or almost clean sulphide mixture with idiomorphic crystals suggests crystallization from a single magmatic solution, and not from several solutions following one another, as was evidently the case for example with the lead-silver-zinc lodes.

5. The deposits often occur in those parts of a gabbro mass which are distinguished by pronounced differentiation of the eruptive rock.

6. The deposits are sometimes crossed by dikes of basic rock, diabase, olivine-diabase, etc., which are to be regarded as later effusions of the same eruption of gabbro. The formation of the ore belongs therefore to the magmatic period of the rock in which it occurs.

7. Some of the deposits are traversed and accompanied by acid leucocratic streaks and dikes which represent the acid segregation products from the gabbro-magma, from which also it follows that the formation of the deposits took place during the magmatic period of the eruptive rock.

8. The characteristic presence of titanomagnetite allows a manner of formation analogous to that of the magmatic titaniferous-iron deposits to be postulated.

9. Pneumatolytic minerals are completely wanting.

10. The minerals usually formed in the wet way are not present as part of the primary formation."

The points in paragraphs 1 and 2 are verified by our work. The ores are so closely connected with a particular type of basic rock

⁶⁸ Ore Deposits: Beyschlag, Vogt, and Krusch; Translated by S. J. Truscott; pp. 286-287. We have revised the translation slightly for the sake of accuracy.

that a genetic relation cannot be doubted. The majority of the deposits are so similar that one method of ore formation alone is involved. The various complicated sets of minerals produced by hydrothermal solutions are either lacking or are developed after the formation of the ores, and hence a close relation to rock crystallization is suggested.

Paragraph 3 has little meaning. Variations in amount of replacement, from incomplete to complete, are found in most ores formed by this process.⁶⁹ The same relations obtain in magmatic ores.

In regard to paragraph 4, Vogt is in error as to the facts. Apparently he had not studied polished ore surfaces, nor has he grasped the meaning of the data set forth in the articles by Dickson⁷⁰ and by Campbell and Knight.⁷¹ The ores of the magmatic stage are introduced into the completely crystallized rock, one after the other, in the following order: (1) magnetite and ilmenite, (2) pyrrhotite, (3) pentlandite, and (4) chalcopyrite. There is positive evidence of a certain amount of the replacement of the earlier ore-minerals by those introduced at later stages. This replacement cannot be explained by corrosion, for there are no intermediate reaction products. The replaced minerals are completely removed by the same agencies that brought in the ores. Vogt recognizes that some of the chalcopyrite is later than the other ore-minerals. He states:⁷² "Die häufig beobachtete Anreicherung des Kupferkieses, namentlich in den peripheren Teilen der Nickel-Magnetkies-lagerstätten muss auf eine besondere magmatische Differentiation innerhalb der Sulfidmagmen beruhen."

To apply his ideas to the facts now established, he would have to postulate a triple or quadruple differentiation within the sulfid magma.

The points in paragraphs 5, 6, and 7 are verified and emphasized by us. Magmatic differentiation is characteristic of the ore-bearing norite, and is especially marked in the vicinity of the ores. Extensive early differentiation calls for an equivalent development of the complementary later differentiation. The ores are a phase of the latter. They are extracted from the magma basin by mineralizers, and are brought to the margin of the deposit and deposited without the development of secondary silicates.

Point 8 is valid.

⁶⁹ Knight (loc. cit.) emphasizes this in his recent discussion of the Sudbury ores.

⁷⁰ Loc. cit.

⁷¹ Loc. cit.

⁷² Beyschlag, Krusch, Vogt.—Die Lagerstätten der nutzbaren Mineralien und Gesteine. Band 1, 284-285 (1910).

Point 9 is true in general, altho tourmaline and garnet are occasionally present and hornblendization generally precedes ore deposition. Vogt evidently assumes that high-temperature deposits formed by mineralizers must be accompanied by destructive pneumatolysis. In this regard it is interesting to note that Vogt's early studies led him to suggest a "combination magmatic segregation with pneumatolysis, according to which metalliferous vapors evolved from the magmas in one place were decomposed in another, depositing the ore in rock already consolidated."⁷⁸ It is needless to state that his early concept approached more closely to the truth than his present hypothesis.

Paragraph 10 may be more accurately stated as follows: The secondary silicates usually formed by destructive pneumatolysis and hydrothermal action are not developed in the magmatic stage of ore formation.

SUMMARY

We conclude with Vogt that the ores are so closely related to the intrusive rocks which contain them, and to the processes of rock differentiation, and differ so markedly from deposits formed by pneumatolytic and hydrothermal processes, that they should be classed as magmatic. However, they cannot be considered as segregated in the molten stage by a "liquation" process, and at a later date intruded into the silicates, for the following reasons:

1. The ore-minerals are formed in an orderly sequence, one after the other. A succession of sulfid differentiations and intrusions is beyond the realm of probability.
2. The ores replace the country rock without reaction rims, and the silicates thus replaced have been completely removed.
3. The minerals are introduced after the magmatic alteration of pyroxene to hornblende, but probably prior to the intense uralitization certain of the deposits have undergone.
4. The temperature of the late stages of differentiation in the presence of mineralizers, and the formation of pegmatite and aplite dikes, is far below the temperature of molten ores, as we know them as furnace products.
5. The phenomena of ore formation and rock replacement are similar in all respects to that of sulfid ore deposition from hydrothermal solutions, except for the comparative absence of secondary silica and silicates. This is a characteristic feature of this type of ore, and is due to the conditions of chemical equilibria under which they are formed.

⁷⁸ Truscott trans. Loc. cit., 289.

BIBLIOGRAPHY OF THE NORWEGIAN DEPOSITS

Vogt, J. H. L.—Bildung von Erzlagerstätten durch Differentiationsprocesse in basischen Eruptivmagmata. *Zeit. f. prakt. Geol.*, Jahrgang 1893, pp. 4-11, 125-143, 257-284.

See also Stelzner-Bergeat, *Die Erzlagerstätten*, I, 48, 1904, for literature of the Norwegian occurrences.

OTHER EUROPEAN OCCURRENCES

SWEDEN

The nickel- and copper-bearing pyrrhotite deposits of Sweden are similar to the more extensive group in Norway. They appear, in general, to have suffered greater alteration than the latter. The largest deposit is at Klefva, in Småland. The ores occur in a quartz norite which has suffered intense uralitization. Some deposits at Kuso and Stattberg occur in mafic (diabase) dikes, and are somewhat similar to the occurrence at Sohland mentioned below.⁷⁴

BADEN, HORBACH, AND TODTMOOS, GERMANY

In the southern Black Forest pyrrhotitic ores occur in small amount in basic inclusions and dikes, extensively altered to amphibolite and serpentine, which occur in granite and gneiss. They are cut by granite-aplite dikes.⁷⁵

SOHLAND AND SWEIDERICH ON THE BOUNDARY BETWEEN SAXONY AND BOHEMIA

Nickel- and copper-bearing pyrrhotite⁷⁶ impregnates the footwall of irregular "basic" dikes which are varied in composition on account of pronounced differentiation.

The ores, according to Beck, surround and cut corroded augite, hornblende, and biotite, and shatter and impregnate the latter. Beck was

⁷⁴ Literature is cited, Beyschlag, Krusch, Vogt, I, p. 294, and Stelzner-Bergeat, I, p. 48.

⁷⁵ Weinschenk, E.—*Die Nickelmagnetkieslagerstätten in Bezirk St. Blasien*. *Zeit. f. prakt. Geol.*, Jahrg. 1907, pp. 73-86. Other literature cited in the textbooks of Beck and Stelzner-Bergeat.

⁷⁶ Beck, R.—*Die Nickelerzlagerstätten von Sohland a. d. Spree und ihre Gesteine*. *Zeit. deutsch. geol. Ges.*, Jahrg. 1903, pp. 296-331. (Includes bibliography of the older literature.)

—*Lehre von Erzlagerstätten*, I, 81-86 (1909).

von Foullon, H. B. *Über einige Nickelvorkommen*. *Jahrb. k.k. geol. Reichsanst.* 302 pp. Wein. (1892).

unable to distinguish pentlandite from pyrrhotite, but we found no difficulty in showing its existence in polished surfaces (see figs. 50 and 51, plate XII).

Beck has proved conclusively that the ores are later than the primary silicates, and believes that they are formed by a "post-volcanic, pneumatolytic phase of rock building."⁷⁷

We have a small suite of specimens from Sohland which include both the unaltered rock and ores. These we have examined in thin sections, and the ores also in polished sections. The unaltered rock is a hornblende-d diabase (called proterobase by Beck), containing plagioclase, augite, brown hornblende, biotite, apatite, titanite, and magnetite. The only alteration products present are a little calcite, tremolite, and chlorite. The hornblende occurs in parallel position with the augite and is evidently a late magmatic mineral.

In thin sections of the ore the only original silicate mineral noted is the hornblende. The feldspars and biotite are altered almost beyond recognition. The alteration products are chlorite, sericite, and uralite. Of especial interest is the uralite, because its relation to the ore-minerals can be definitely established.⁷⁸

Fig. 50 shows the general relations of the minerals in the polished sections. A study of our sections together with Beck's original figures (especially his fig. 4 of plate XIII) makes it almost certain that the ore-minerals were formed later than the hornblende, but earlier than the uralite. Uralite develops on the ends of the hornblende prisms in parallel position, and occasionally completely replaces the hornblende. Frequently the uralite needles project out into the ore-minerals, as in fig. 51, and in Beck's fig. 4. This affords conclusive evidence of post-mineral alteration. For this reason we believe that the Sohland ore is magmatic in spite of the extensive alteration. The relation of the ore-minerals to the silicates was evidently established during the magmatic period, and not modified by later hydrothermal action.

⁷⁷ Quoted by Berg, *opus cit.*, 108.

⁷⁸ In the literature the term uralite has been used in two senses, (1) for hornblende rims around pyroxene, (2) for the fine fibrous aggregates of tremolite (including actinolite). The first usage we have avoided entirely, and in the few cases in which the terms uralite and uralitization are used by us, these are to be given the second meaning. The hornblende rims appear to be late magmatic and the tremolite a post-magmatic, hydrothermal mineral.

GROUP II. MAGMATIC CHALCOPYRITE-BORNITE DEPOSITS

OOKIEP, NAMAQUALAND, SOUTH AFRICA

GEOLOGY

The important mines of this region occur as great lenticular deposits in the vicinity of the town of Ookiep, Namaqualand, Cape Colony, South Africa. They lie about sixty miles east from the Atlantic coast, and about twice as far south of the Orange river.

The ore bodies occur as shoots or lenses, and again as disseminated sulfid particles, in a system of dikes and sills, which give evidence of remarkably extensive differentiation. Rogers⁷⁸ describes the "ore bearer" as including rock varieties made up almost wholly of any one of the following minerals: magnetite, quartz, plagioclase, hypersthene, hornblende, and biotite, and also intermediate varieties such as norite, mica diorite, augite diorite, and diorite. A single dike may show one or more varieties of rock, and where several kinds occur in one intrusive the contacts are sharp, suggesting differentiation before intrusion.

According to Kuntz⁷⁹ the intrusive rock occurs as dikes of great extent, or again as detached stocks. To these Rogers adds: pipes, nearly horizontal sheets, and branching bodies; and states that 344 intrusive bodies have been mapped up to date. These dikes cut the "fundamental gneiss" of South Africa, and the ore occurs as shoots, occupying a portion of the entire width of the dikes, and also penetrates the gneiss adjacent to the dikes.

The principal ore-minerals are given as magnetite, pyrrhotite, bornite, and chalcopryite. We find that hematite is also an important constituent of the ores.

The chief mines are Ookiep, Specktakel, Nababeep, and Ookiep East, all situated near the town of Ookiep, and the Tweefontein mine is located on a dike to the north. The ore shoots are lenticular masses of high grade ore with greatest dimension in a horizontal direction. For example, the Ookiep ore-body, according to Ronaldson,⁸⁰ is 1300 feet

⁷⁸ Rogers, A. W.—The nature of the copper deposits of Little Namaqualand, *Proceed. Geol. Soc. S. Africa*, 1916, pp. xxi-xxxix.

⁷⁹ Kuntz, J.—Copper ore in southwest Africa. *Trans. Geol. Soc. South Africa*, 7, 70-76 (1904).

⁸⁰ Ronaldson, J. H.—Notes on the copper deposits of Little Namaqualand. *Trans. Geol. Soc. S. Africa*, 8, 158-167 (1905).

wide and 300 feet deep. The Tweefontein mine contains a series of three lenses, one under the other.

On account of the occurrence of the copper ore intimately mixed with apparently unaltered constituents of the dikes, a magmatic origin was early assigned to these deposits. In 1857, Wiley⁸¹ stated that the ores were of magmatic origin, and Daintre in 1878 makes the plain statement that "the ores are of magmatic origin, occurring in unaltered feldspathic dikes from specks the size of a pin point to many tons."⁸²

MICROSCOPIC DESCRIPTIONS

We are indebted to Dr. A. W. Rogers, Director of the Geological Survey of the Union of South Africa, for an excellent suite of specimens of ores and accompanying rocks from the Ookiep mines. These rocks vary from an almost pure hypersthene rock (hypersthenite), through norite and mica diorite, to an almost pure plagioclase rock (anorthosite). The specimens include both lean and rich norite and mica diorite, but the anorthosite is almost free from the ore-minerals, and the hypersthenite carries a fair amount of the ore-minerals.

Ore-bearing Hypersthenite from the Nababeep Mine.—The general relations of the ore-minerals to the silicates of the hypersthenite are well shown in fig. 54. The ore-minerals occur in anhedral which are often elongate and pass into vein-like forms. It is probable that bornite has formed by the replacement of the magnetite, for they show the same general shape.

Fig. 52 represents a thin section of the hypersthenite. The transparent mineral is hypersthene. The black area is mainly magnetite, but a little bornite is also present, and in some places in the section bornite predominates over magnetite. The ore-minerals surround the hypersthene crystals, and in one place (just below the center) a veinlet of magnetite cuts across a hypersthene crystal.

Besides hypersthene the only original silicates present are biotite and plagioclase, which occur in minor amounts. The plagioclase is shown in little patches in fig. 53. The association of ores with these areas of feldspar, which considered with respect to the enclosing rock are of the nature of "acid extracts," is significant. The feldspar seems to be more readily replaced by the ore-minerals than the hypersthene. Note the small magnetite euhedra within the hypersthene.

⁸¹ Wiley, A.—Report on the mineral and geological structure of South Namaqualand. Parliamentary Report, Cape Town, no. 36, p. 30 (1857).

⁸² Quar. Jour. Geol. Soc., 34, 434 (1878).

PETROGRAPHY AND MINERALOGY

The ore-bearing rock may be classed broadly as norite (\pm quartz, olivine, and diallage), and in a few localities it is so extensively uralitized as to be classified by some authorities as a "gabbro-diorite." Each individual stock has undergone extensive differentiation, especially in the vicinity of the ore bodies. The basic segregations are mixtures of bronzite, hornblende, olivine, and biotite. They occur in masses, smaller inclusions, and, at Romsaas, as a peculiar orbicular norite. The felsic




Fig. 3. Erteli mine, Ringerike, Norway. (After Vogt, *Zeit. f. prakt. Geol.*, fig. 8, plate V, Jahrgang 1893.)

products of differentiation are as prominent as the mafic, and occur as masses of granite, streaks and dikes of pegmatitic and aplitic material, and small marginal veinlets of aplite and pegmatite. A similar differentiation appears to be a characteristic of all deposits of this class. The margins of the stocks often show severe fracturing and brecciation, and the ore may cement these igneous breccias, as at Sudbury.

According to our ideas, it would appear that (1) the "basic" portions of the norite solidified by the sinking of crystals; (2) portions of this "basic" material were caught up as inclusions in the normal norite, especially along the margins of the stocks; (3) the main mass of the rock crystallized as norite; (4) felsic dikes and masses were squeezed out as marginal segregates during a period of peripheral brecciation, which facilitated the concentration and escape of mineralizers from the deeper portions of the magma; (5) ore deposition, at a late stage, was preceded, accompanied, and followed, by the "acid" secretions, as the ore is found in the felsic dikes, is later than the silicates of the same, and the main ore masses are cut by the dikes.




Fig. 4. Detailed cross-section of ore-body at the Meinkjær mine, Norway. (After Vogt, *Zeit. f. prakt. Geol.*, fig. 29, p. 136, Jahrgang 1893.)

The ores occur chiefly at the margins of the stocks, and to a minor amount as segregations within the norite, and also as impregnations in the schists and gneisses in which the norite masses are enclosed. As has been mentioned, the ores are further localized where marked differen-



Fig. 5. Pyrrhotite gabbro from Erteli mine, Norway. Pyrrhotite veinlet cutting feldspar and diorite with rim of garnet. (x 100.) (After Vogt, *loc. cit.*, p. 139.)



Fig. 6. Pyrrhotite veinlets cutting hornblende. Flaad mine, Evje, Norway. (x 50.) (After Vogt, *loc. cit.*, p. 139.)

tiation has taken place. At the Flaad mine, especially, the ores are closely related to felsic differentiates.

The ores show the same group of minerals (chiefly magnetite, pyrrhotite, pentlandite, and chalcopyrite), and the same relation to each other and to the schists, as we have found in the deposits studied elsewhere. Pyrite is mentioned by Vogt as of common occurrence in these

biotite-rich segregation in the diorite. The biotite, itself probably a late magmatic mineral, has been replaced along its cleavage planes by chalcopyrite. This produces a structure which can be distinguished from that developed by the replacement of the sulfids by later hydrothermal minerals such as sericite, chlorite, and tremolite. (See for examples figs. 82 and 83, plate XX.)

SUMMARY

The Ookiep ores are of importance in establishing the type of magmatic copper deposits. All who have studied these ores have classified them as magmatic ores, but in spite of this they have received scant attention by the authors of textbooks and treatises on ore deposits. The principal reason for the failure to give these deposits full standing in the magmatic class is the presence of bornite. The Ookiep ores are magmatic, if such exist. They are the least altered of any we have examined in this study. The ore-bearing (chalcopyrite-bornite) norite from the Tweefontein mine is much freer from alteration than the average igneous rock. It is true that the mica diorite and hyperstheneite at Ookiep locally show some alteration, but the significant thing is that the ore-minerals in the altered and unaltered rocks show exactly the same relation to the silicate minerals.

BIBLIOGRAPHY OF THE OOKIEP DEPOSITS

- Delesse, M.—Sur les mines de cuivre du Cap de Bonne Esperance. *Ann. des Mines*, 5me serie, 8, 186-212 (1855).
- Wiley, A.—Report on the mineral and geological structure of South Namaqualand. Parliamentary Report, Cape Town, no. 36, p. 30 (1857).
- Zerrener, C.—Reise des Ingenieurs A. Thiers nach den Kupfer Bergwerken Namaqualands in Sud Africa. *Berg. und Hüttenm. Zeitung* 1860, pp. 41-44 and 53-54.
- Knopf, A.—Über die Kupfererzlagerstätten von Klein Namaqualand und Damara-land. *Neues Jahrb. f. Min. Geol. u. Pal.*, 1861, pp. 513-550.
- Schenk, A.—Die Kupfererzlagerstätten von Ookiep in Klein Namaqualand. *Zeit. de deutsch. geol. Ges. Verhand. der Ges.*, 53, 64-65 (1902).
- Kuntz, J.—Copper ore in Southwest Africa. *Trans. Geol. Soc. South Africa*, 7, 70-76 (1904).
- Kupfererzsvorkommen in Südwestafrika. *Zeit. f. prakt. Geol.*, 12, 199-202 (1904).
- Ronaldson, J. H.—Notes on the copper deposits of Little Namaqualand. *Trans. Geol. Soc. South Africa*, 8, 158-167 (1905).
- Stutzer, O.—Magmatische Ausscheidungen von Bornit. *Zeit. f. prakt. Geol.*, 15, 371 (1907).
- Rogers, A. W.—The nature of the copper deposits of Little Namaqualand. *Proceed. Geol. Soc. S. Africa*, 1916, pp. xxi-xxxiv.

ENGELS MINE, PLUMAS COUNTY, CALIFORNIA

GEOLOGY

A rather unique copper deposit, which we believe to be magmatic, is now being mined by the Engels Copper Mining Company in the northern part of Plumas county, California. This deposit has been studied and described by Turner and Rogers.⁸⁸

The ore occurs in a remarkably fresh norite-diorite occurring at the extreme northern end of the great Sierra Nevada batholith of granodiorite. There is no evidence of dynamic metamorphism and none of contact metamorphism, for the older rocks into which the norite-diorite is intrusive are nearly five miles distant. The main ore body is a tabular ore shoot which has a nearly vertical attitude. It is difficult to see how this ore body could possibly be explained by gravitative adjustment due to the sinking of sulfids in the molten magma. Narrow pegmatite dikes are occasionally found cutting the norite-diorite.

MICROSCOPIC DESCRIPTIONS

The predominant igneous rock at the Engels mine is a norite-diorite with about 46 per cent of silica. The principal minerals are plagioclase (andesine-labradorite, Ab_1 An_1), hypersthene, diopside, hornblende, and biotite. The norite-diorite varies from a rather fine-grained hypersthene-plagioclase rock to a coarse-grained hornblende-plagioclase rock. One of the typical fine-grained rocks (from no. 2 level) is illustrated in fig. 72. The opaque minerals here are magnetite and hematite. They surround and replace the silicates, especially hypersthene and biotite. Evidence of replacement is shown at many spots in the photomicrograph. A magnified view of one of these is shown in fig. 74. The magnetite and hematite have surrounded hypersthene and plagioclase, and have penetrated and replaced a biotite crystal. It will be noticed in fig. 72 that the magnetite occurs in both euhedral and anhedral crystals. The euhedral crystals occur within the silicate minerals, and the anhedral crystals mainly along the boundaries of adjacent silicate anhedral. This is also well shown in fig. 73. In this figure there is a magnetite crystal which shows anhedral development on the boundary between the hypersthene and plagioclase, and euhedral development within the hypersthene. This is an argument in favor of the late magmatic origin of

⁸⁸ A geologic and microscopic study of a magmatic copper sulphide deposit in Plumas county, California, and its modification by ascending secondary enrichment. *Econ. Geol.*, 9, 359-391 (1914).

PETROGRAPHY AND MINERALOGY

The ore-bearing rock may be classed broadly as norite (\pm quartz, olivine, and diallage), and in a few localities it is so extensively uralitized as to be classified by some authorities as a "gabbro-diorite." Each individual stock has undergone extensive differentiation, especially in the vicinity of the ore bodies. The basic segregations are mixtures of bronzite, hornblende, olivine, and biotite. They occur in masses, smaller inclusions, and, at Romsaas, as a peculiar orbicular norite. The felsic




Fig. 3. Erteli mine, Ringerike, Norway. (After Vogt, *Zeit. f. prakt. Geol.*, fig. 8, plate V, Jahrgang 1893.)

products of differentiation are as prominent as the mafic, and occur as masses of granite, streaks and dikes of pegmatitic and aplitic material, and small marginal veinlets of aplite and pegmatite. A similar differentiation appears to be a characteristic of all deposits of this class. The margins of the stocks often show severe fracturing and brecciation, and the ore may cement these igneous breccias, as at Sudbury.

According to our ideas, it would appear that (1) the "basic" portions of the norite solidified by the sinking of crystals; (2) portions of this "basic" material were caught up as inclusions in the normal norite, especially along the margins of the stocks; (3) the main mass of the rock crystallized as norite; (4) felsic dikes and masses were squeezed out as marginal segregates during a period of peripheral brecciation, which facilitated the concentration and escape of mineralizers from the deeper portions of the magma; (5) ore deposition, at a late stage, was preceded, accompanied, and followed, by the "acid" secretions, as the ore is found in the felsic dikes, is later than the silicates of the same, and the main ore masses are cut by the dikes.

The ores occur chiefly at the margins of the stocks, and to a minor amount as segregations within the norite, and also as impregnations in the schists and gneisses in which the norite masses are enclosed. As has been mentioned, the ores are further localized where marked differen-




Fig. 4. Detailed cross-section of ore-body at the Meinkjær mine, Norway. (After Vogt, *Zeit. f. prakt. Geol.*, fig. 29, p. 136, Jahrgang 1893.)



Fig. 5. Pyrrhotite gabbro from Erteli mine, Norway. Pyrrhotite veinlet cutting feldspar and diorite with rim of garnet. ($\times 100$.) (After Vogt, *loc. cit.*, p. 139.)



Fig. 6. Pyrrhotite veinlets cutting hornblende. Flaad mine, Evje, Norway. ($\times 50$.) (After Vogt, *loc. cit.*, p. 139.)

tiation has taken place. At the Flaad mine, especially, the ores are closely related to felsic differentiates.

The ores show the same group of minerals (chiefly magnetite, pyrrhotite, pentlandite, and chalcopyrite), and the same relation to each other and to the schists, as we have found in the deposits studied elsewhere. Pyrite is mentioned by Vogt as of common occurrence in these

SUMMARY

The Engels mine, which is rapidly becoming one of the important copper mines of California, is unique in that magmatic bornite is the principal ore-mineral. The ore occurs as a vertical ore shoot, and in a massive norite-diorite, in such relation to the enclosing rock that only a magmatic origin seems probable. Bornite and associated chalcopyrite are formed by mineralizers at a late magmatic stage, and not by hydrothermal solutions. This is proved by the comparative freedom from alteration in many of the ores and associated rocks. The silicification accompanying hydrothermal deposits generally is practically absent. Locally there may be sericitization and chloritization, but the bornite and chalcopyrite are no more abundant in the ores thus affected than in the unaltered ores. The secondary copper minerals, covellite, chalcocite, and chalcopyrite of the second generation, have developed at a later stage, in part by hypogene and in part by supergene solutions.

BIBLIOGRAPHY OF THE ENGELS ORE DEPOSIT

- Aubury, L. E.—The copper resources of California. Bull. no. 50, California State Mining Bureau, pp. 185-186 (1908).
- Turner, H. W., and Rogers, A. F.—A geologic and microscopic study of a magmatic copper sulphide deposit in Plumas county, California, and its modification by ascending secondary enrichment. *Econ. Geol.*, 9, 359-391 (1914).
- Read, T. T.—The Engels mine and mill. *Min. and Sci. Press*, 111, 167-171 (1915).

REMARKS ON CERTAIN OTHER DEPOSITS THAT HAVE
BEEN CLASSIFIED AS MAGMATIC

PYRITIC DEPOSITS

Beyschlag, Krusch, and Vogt⁸⁷ classify certain pyrite ores as magmatic ores under the heading "Die intrusiven Kieslagerstätten." Among these they place the Rio Tinto, Bodenmais, Sain Bel and Chessy, Agordo, and numerous Norwegian deposits. It is certain that some of these pyritic ores are hydrothermal in origin. The Rio Tinto deposits, for example, are due to metasomatic replacement of crushed and sheared zones by hydrothermal solutions, as was definitely proved by Finlayson.⁸⁸ The Rio Tinto and other pyritic deposits, such as the Rammelsberg, are placed by Lindgren⁸⁹ in his division "Deposits formed at Intermediate Depths." Lindgren, however, recognizes that some of the pyritic deposits may be due to the injection of molten sulfids.

The so-called intrusive or injected pyritic ores occur for the most part in gneisses. Igneous rocks are often closely associated, tho not always. The igneous rocks, however, are altered rocks, such as saussurite-gabbro. The ore-minerals are associated with typical metamorphic minerals such as cordierite, andalusite, anthophyllite, and chloritoid.

As these ores occur in areas of regional metamorphism, and are usually found in gneisses, schists, and not often in the igneous rocks themselves, it is difficult to see why they are placed with the magmatic ores. It seems far more reasonable to classify them as metamorphic ores, as Berg⁹⁰ does. We have not had an opportunity of examining many of these ores, but certain facts derived from our study of the undoubted magmatic ores leads to the conclusion that these are not of this type, or at least that the burden of proof rests upon any one who classifies them as such.

In the first place, we seriously doubt whether pyrite is a characteristic magmatic mineral. In our present study we have not found pyrite in any of the typical disseminated magmatic ores, and our speci-

⁸⁷ Loc. cit., 1, 298 (1910).

Ore Deposits. English translation by Truscott, 1, 301 (1914).

⁸⁸ Finlayson, A. M.—The pyritic deposits of Huelva, Spain. *Econ. Geol.*, 5, 357-372, 403-437 (1910).

⁸⁹ Mineral Deposits, 602 et seq. (1913).

⁹⁰ Loc. cit., 114.

mens include representative suites of most of the important deposits except those from Norway. It is true that pyrite occurs in some of the Sudbury mines; we have examined several of these occurrences, and have found the pyrite to be distinctly later than the nickel-copper sulfids.

Later mineralization seems to be more prominent at the Worthington mine than at any other locality in the Sudbury district. Walker⁹¹ says: "It is difficult to avoid the conclusion that the ores as they are now found at the Worthington mine have been subject to rearrangement by aqueous agencies since the solidification of the rock and sulfids from the original magma." Pyrite is evidently not a typical mineral at Sudbury. Browne,⁹² for example, says: "Practically speaking, there is no pyrite, marcasite, or any other sulphide (he has previously mentioned pyrrhotite, chalcopyrite, and pentlandite) found in the great ore bodies."

Pyrite is entirely absent in the ores from the Engels mine. This might be attributed to lack of sufficient iron to combine with sulfur; but in the Namaqualand ores there was sufficient iron to form pyrrhotite, yet pyrite is entirely absent.

As a matter of fact, pyrrhotite seems to take the place of pyrite in the magmatic ores. The work carried on⁹³ at the Geophysical Laboratory, proving that pyrrhotite is the iron sulfid stable at high temperature, may explain this. Pyrite has usually been considered a persistent mineral.⁹⁴ Its maximum development, however, is reached in hydrothermal deposits, and it is less important in contact deposits and in those formed at low temperatures by meteoric water. True, pyrite is common in igneous rocks; but in the great majority of cases it has been introduced by hydrothermal solutions. There are a few authentic cases⁹⁵ of pyrite in fresh unaltered igneous rocks, and so its occurrence as a magmatic mineral can not be absolutely denied; but it seems certain that it is of minor importance as a magmatic mineral.

⁹¹ Walker, T. L.—Certain mineral occurrences in the Worthington mine, Sudbury, Ontario, and their significance. *Econ. Geol.*, 10, 542 (1915).

⁹² Browne, D. H.—Notes on the origin of the Sudbury ores. *Econ. Geol.*, 7, 468 (1906).

⁹³ Allen, E. T., Crenshaw, J. L., and Johnston, J.—The mineral sulphides of iron. *Am. Jour. Sci.* (4), 33, 169-236 (1912).

⁹⁴ Lindgren, W.—The relation of ore-deposition to physical conditions. *Econ. Geol.*, 2, (1907).

⁹⁵ Lindgren, W.—The gold-quartz veins of Nevada City and Grass Valley districts, California. 17th Ann. Rept. U. S. Geol. Surv., pt. II, 66 (1896).

Spurr, J. E., and Garrey, G. H.—Prof. Paper no. 63, U. S. Geol. Surv., 388 (1908).

MAGNETITE-ILMENITE DEPOSITS

Geologists apparently find little difficulty in accepting the titaniferous magnetites as magmatic deposits, probably because magnetite is a common accessory mineral of igneous rocks, but appear to be more cautious in regard to the sulfid deposits. We have examined a number of examples of the magmatic iron ores in "basic" rocks, and find that the relations of the ores to silicates are similar in all respects to those of the sulfid deposits. In the latter the sulfids may fail locally and the ores become identical with the low-grade titaniferous iron ores. The sulfid rich and sulfid free magnetite deposits alike develop subhedral crystals within the silicates; and the larger anhedral crystals cut, surround, and replace the silicates. They are often unaccompanied by secondary alteration products, and where these are present, they are later than the ores.

In order to check our incomplete study, we have examined the literature of these deposits, and find, contrary to the general impression, that the majority of those who have made a careful microscopic study of the ores conclude that they are later than the silicates. As a result of a similar review of the literature, Lindgren⁸⁶ states: "In these differentiated magmas ilmenite and magnetite have, as a rule, crystallized after the silicates." Berg⁸⁷ notes: "Die grosseren Erzmassen sind stets Anhaufungen des jungeren Erzes, umschliessen also einzelne Krystalle von Hypersthen, Diallag, Olivin, und Feldspat."

OTHER MAGMATIC IRON ORES

It is not our purpose to take up the remaining types of iron ores for which a magmatic origin has been suggested, the origin of some of which is obscure on account of the metamorphism they have undergone. In general, those deposits related to the less "basic" rocks, such as the Kiirunavaara type,⁸⁸ and certain of the Adirondack ores,⁸⁹ show evidence of increasing activity of mineralizers, with the occasional development of pneumatolytic and allied minerals, and a tendency to migrate further out from the mother rock.

In general, iron oxids, like the magmatic sulfids, appear to be concentrated, not during the early magmatic stages, but during the later stages and under the influence of mineralizers.

⁸⁶ Mineral Deposits, 749.

⁸⁷ Loc. cit., 102.

⁸⁸ Igneous rocks and iron ores of Kiirunavaara, Luossavaara, and Tuollavaara. Econ. Geol., 5, 696-718 (1910).

⁸⁹ Newland, D. H.—On the association and origin of the non-titaniferous magnetites in the Adirondack region, 2, 763-773 (1907).

CHROMITE DEPOSITS

Of the remaining ore-minerals concentrated as magmatic segregations, nickel-iron, gold, and platinum, do not occur in sufficient amounts to be considered as ores.

Chromite in unaltered rock from Norway, has been described by Vogt,¹⁰⁰ and is considered by him to be the oldest mineral of the rock, on account of its occurrence as sharp octahedra. Examination of his figures shows irregular as well as euhedral forms; and the alternate hypothesis that the chromite octahedra are formed at a late stage, is worthy of consideration.

Vogt states: "The chromite deposits in peridotite show accordingly the same geological, petrographical and morphological characteristics as those of titaniferous iron in gabbro, and the general genetic statements afterwards enumerated in connection with the titaniferous iron deposits hold good also for the occurrence of chromite."¹⁰¹

¹⁰⁰ Beiträge zur genetischen Classification der durch magmatische Differentiationsprocesse und durch Pneumatolyse entstanden Erzvorkommen. Zeit. f. prakt. Geol., Jahrg. 1894, pp. 384-393.

¹⁰¹ Ore Deposits: Beyschlag, Vogt, and Krusch; Translated by Truscott, (1914), 246.

PART III. SUMMARY AND CONCLUSIONS

CRITERIA FOR THE RECOGNITION OF MAGMATIC ORES

Any discussion of the criteria by which the magmatic ores may be separated from other types of ores must be somewhat tentative in nature, as long as the latter have not received detailed comparative study.

The fact that the ores migrate only a short distance, if at all, into the adjoining rock, distinguishes them from contact deposits formed chiefly in the intruded rock, and from many other high-temperature deposits.

Comparison, then, must be made with ores that occur in, and as an integral part of, igneous rocks.

The fact that the only high-temperature alteration mineral present in appreciable amount is hornblende, which we believe is formed by magmatic processes, and that tourmaline and garnet are only occasionally developed, distinguishes these deposits from those in which destructive pneumatolytic and contact action is prominent.

The fact that hydrothermal alteration of the silicates is often minor in amount, is invariably later than the ores, and generally is not accompanied by the deposition or migration of ores, differentiates these ores from those of hydrothermal origin.

The absence of silicification and albitization, and the only occasional development of considerable amounts of sericite, separate these deposits from moderate- and high-temperature ores in igneous rocks, such as the copper ores of Bingham and Butte and the gold ores at Treadwell, Alaska, etc.

There is definite order of succession of the magmatic ore-minerals, as follows: magnetite, hematite, pyrrhotite, pentlandite, chalcopyrite, bornite. Any change in this order is due to rearrangement subsequent to the magmatic period.

Pyrite is not a typical magmatic mineral; if present in a magmatic ore, it is introduced later than the magmatic period.

SUMMARY OF THE CHARACTERISTICS OF MAGMATIC ORES

Contrary to general opinion, the magmatic sulfids are formed by replacement of the silicates after the solidification of the igneous rock. Notwithstanding this, we retain the term "magmatic ore deposits" for the types of ores described in this paper, because they have a close genetic relation to the intrusive rock in which they occur, and because they are formed within the magmatic stage as defined by us. Regardless of any theory as to their genesis, however, these deposits have definite and easily recognizable characteristics, which distinguish them from all other types of ore deposits.

The characteristics of the magmatic sulfid ores as brought out by our study may be summarized as follows:

(1) They occur in subsilicic rocks of the norite, gabbro, peridotite, or related families.

(2) In most occurrences the containing rock is either dominantly subsilicic, with minor amounts of complementary persilicic differentiates (Insizwa), or occurs as lenses of mafic rock in a large granitic intrusion (Golden Curry).

(3) The subsilicic rock occurs generally as small dikes, sills, or stocks, and rarely as a large laccolith (Sudbury).

(4) In most cases the ore-bearing rock has undergone marked differentiation, and the differentiated portions are sharply separated and do not grade into each other (Ookiep).

(5) The ore may occur in any variety of rock produced by differentiation, but in any one locality the ore shows marked preference for certain types of rock and occurs sparingly in others (Ookiep).

(6) Pegmatite and aplite dikes often cut, and are therefore later than, the ore bodies (Erteli).

(7) The ore is generally segregated at the margins of the intrusives, but occasionally occurs as lenses or tabular ore shoots well within the intrusive magma (Engels). In sills, the ore is usually at the base of the intrusives (Insizwa); in dikes, it often is formed along the footwall (Sohland), or as columnar or irregular shoots occupying the entire width of the dike (Copper Cliff).

(8) Very often the ore is concentrated in those portions of the intrusive which have suffered brecciation during intrusion (Sudbury).

(9) The ore migrates only a short distance into the adjacent rock, from a few inches to a few score of feet at most (Sudbury).

(10) The "offset" deposits, formed in dikes at some distance from the main intrusive, are accompanied and cut by veins carrying ore and gangue minerals of hydrothermal origin (Sudbury).

(11) There is no essential distinction between the sulfid group and the magnetite-ilmenite group as to the origin or the relation of the ores to the silicate minerals. In all the magmatic ores examined, ore deposition takes place at the close of the magmatic period.

(12) There are two general classes of the magmatic sulfid ores: (a) pyrrhotite-chalcopyrite deposits, and (b) chalcopyrite-bornite deposits.

(13) The so-called pyritic intrusive ores are not magmatic.

(14) The principal ore-minerals of the magmatic period include magnetite, ilmenite, hematite, pyrrhotite, pentlandite, chalcopyrite, and bornite.

(15) Pyrrhotite and bornite have not been found together in magmatic ores.

(16) Pyrite is not a typical magmatic mineral.

(17) The ore-minerals are formed at a late magmatic stage by a partial replacement of the silicate minerals. The ores surround, embay, and cut all the earlier silicates. They penetrate the cleavable minerals. They occasionally occur as sharp veinlets which lead out from the larger sulfid masses. Selective replacement is shown by the preservation in the ores of an original graphic texture of the rock.

(18) There is a definite order of formation of the principal magmatic ore-minerals. This order is as follows: magnetite-ilmenite (intergrowth), hematite, pyrrhotite, pentlandite, chalcopyrite, and bornite.

(19) There is evidence of the replacement of one magmatic ore-mineral by another.

(20) Euhedral magnetite and probably other minor accessories occurring in euhedral crystals, such as apatite, zircon, titanite, etc., are also formed at a late magmatic stage.

(21) There is clear evidence of the magmatic alteration of pyroxene to hornblende prior to the introduction of the ore-minerals.

(22) Hydrothermal alteration, altho seldom lacking in magmatic ores, is relatively insignificant, and is distinctly later than the magmatic ore period. The silicates of the hydrothermal period include tremolite, anthophyllite, sericite, chlorite, and serpentine. These secondary silicates often replace the magmatic ore-minerals in veinlets and in sharp

lath-shaped crystals without causing any change or migration of the ore-minerals.

(23) The attention given these alteration products in order to determine their relative age, may have given an erroneous impression as to their relative abundance. In most of the ores studied they are present only in minor amounts. Many of the magmatic ores are as free from alteration products as the average unmineralized igneous rock.

(24) However, a minor amount of rearrangement, consisting in the production of microscopic crystals of pentlandite and chalcopyrite of the second generation, has been detected in the pyrrhotitic ores. In the chalcopyrite-bornite ores there has been some migration, resulting in the formation of minor amounts of covellite, chalcocite, and chalcopyrite of the second generation. This alteration is only prominent where there has been an abnormal development of sericitization.

(25) The rôle of mineralizers in magmatic differentiation has not been sufficiently emphasized. The crystallization of the early formed minerals in the magma involves the complementary process of the "squeezing out" of the residual fluid. This process is not merely a mechanical one, but is also due to gaseous extraction.

(26) The typical magmatic deposits, unaccompanied by high-temperature alteration, with the exception of magmatic hornblende, are chiefly developed in "basic" rocks. Ore deposits genetically related to persilicic rocks show intense rock alteration, probably the result of mineralizers more "active" than those accompanying the subsilicic rocks.

(27) There is a parallelism between the various groups of high-temperature deposits, of which the magmatic ores are one division. In all groups, high-temperature silicates precede the introduction of the ore, and hydrothermal stages follow. In contrast with the magmatic deposits, the non-magmatic ores are characterized by a complex set of pre-mineral silicates, and the hydrothermal stage is generally the most important period of ore introduction.

(28) The temperature at which the introduction of ore-minerals is initiated is about the same for all the high-temperature deposits; probably not higher than 300-400° C.

(29) Gradations between the typical magmatic ores and other high-temperature deposits are shown by the local development of garnet and tourmaline. Further, the gradations of these into intermediate-temperature and low-temperature deposits is strong evidence of the magmatic origin of ore deposits in general.

(30) The following orderly series of events is recognized in magmatic deposits:

- (a) Crystallization of primary silicates;
- (b) The development of hornblende and biotite, and occasionally tourmaline and garnet, as magmatic alteration products;
- (c) The introduction of the ore-minerals;
- (d) A small amount of rearrangement of the ores and the development of secondary silicates by hydrothermal solutions.

Department of Geology,
Leland Stanford Junior University,
October, 1916

EXPLANATION OF PLATES AND METHODS OF PREPARING PHOTOGRAPHS AND SECTIONS

The photographs are of two types: thin sections and polished sections. The thin sections are necessary for the identification of the silicates; the polished sections for the identification of the ore-minerals,




Fig. 6A. Drawn from a photograph of a polished thin section of ore from the Stobie mine, Sudbury. White areas are silicates. Lined area is opaque band as viewed by transmitted light. Black band is veinlet as shown by reflected light. ($\times 175$)

and to show the relations of the various minerals to each other. In examining the photographs note that the ore-minerals are black and the silicates light in thin sections, while the silicates are dark gray and the ore-minerals white or light gray in the polished sections.

In taking the photographs we have used Wratten panchromatic plates, selecting in each case the color screen best adapted to bring out all the minerals in the polished section.

In the microscopic investigation we examined (1) thin sections with the polarizing microscope; (2) polished sections with the reflecting microscope; and (3) polished thin sections (combination sections) with both transmitted and reflected light.

It is now generally admitted that the examination and determination of the opaque minerals in thin sections is at best unsatisfactory, and that only the grosser relations of the opaque minerals to each other can be made out, and that those details requiring high magnifications are lost entirely.

In addition to the above, we have found that the relations of the opaque minerals, as a group, to the transparent minerals, are shown more accurately in polished surfaces than in thin sections, and that the relations of the transparent silicates to each other are sharper when viewed with the reflecting microscope than with the polarizing microscope, altho the identification of the transparent minerals must be made with the latter.

The reason for this is simple. In examining a thin section, one looks thru a certain thickness of rock, and not at a single definite surface. All contacts between the opaque and transparent minerals that are not perpendicular to the surface of the slide are widened so that the black area represents the projection on the surface of the section of all the opaque material contained in the slide. A thin section does not show, therefore, the relation of the opaque and the transparent minerals at one definite level as do the polished sections. Fig. 6a, page 74, was drawn from a photograph of a polished thin section viewed under both transmitted and reflected light. The ruled band represents the shadow cast by the opaque material as viewed with transmitted light, and the thin black line at the margin of the band shows the true thickness of the veinlet as registered by the light reflected from the polished surface.

In order to take advantage of the reflecting microscope for this work, it was necessary to develop special methods of polishing in order to give a satisfactory brilliant surface to the silicates as well as to the hard and soft opaque minerals. As a matter of fact, the polishing of all the different constituents in a single section of complex ore is an art, and, up to date, only crude methods of polishing have been described in the literature.

Our methods, in brief, consist in grinding on a glass plate a plane surface with the minimum of pits, using successively finer grades of carborundum, and then "sixty-minute" carborundum, and finally a still finer carborundum prepared in our laboratory.

The final polishing is done with fine cambric on rapidly rotating wheels (1700-3400 r.p.m.). Silicates, magnetite-ilmenite, pyrite, etc., are polished by carborundum, prepared by grinding the "sixty-minute" grade two weeks in a ball mill, and then floating off the finest product for this purpose. Chrome oxid is used to polish sulfids of average hardness, and aluminum oxid for the softer sulfids. These powders are applied one after the other, or as mixtures, depending upon the character of the specimen. To develop a polish of the highest brilliancy on the sulfids—for example to bring out pentlandite enclosed in pyrrhotite—aluminum oxid is used on the finest billiard felt.

In most cases we find it satisfactory to use one side of a sawed specimen for the polished section and the adjacent surface for the thin section. In some cases a thin section is given a high polish, which acts like a cover-glass in giving clear images with transmitted light, and also gives brilliant reflections with reflected light.

PLATE I
PHOTOGRAPHS OF POLISHED SPECIMENS OF SUDBURY ORES

FIG. 7

Rich ore (pyrrhotite and pentlandite) from Copper Cliff mine, showing unreplaced silicates [dark areas] of the granitic rock.

Nat. size

FIG. 8

Ore from the Creighton mine showing chalcopyrite veinlets.
x $1\frac{2}{3}$ diameters

FIG. 9

Nodule of "basic" material in granitic rock which has been largely replaced by pyrrhotite [light gray].
Creighton mine.

x 87/100

FIG. 10

Pyrrhotite [light gray] impregnating and replacing "greenstone schist." Garson mine.

x 8/10




Fig. 7




Fig. 8




Fig. 9




Fig. 10

PLATE I
PHOTOGRAPHS OF POLISHED SPECIMENS OF SUDBURY ORES

FIG. 7

Rich ore (pyrrhotite and pentlandite) from Copper Cliff mine, showing unreplaced silicates [dark areas] of the granitic rock.

Nat. size

FIG. 8

Ore from the Creighton mine showing chalcopyrite veinlets.
x 123 diameters

FIG. 9

Nodule of "basic" material in granitic rock which has been largely replaced by pyrrhotite [light gray] Creighton mine.

x 87/100

FIG. 10

Pyrrhotite [light gray] impregnating and replacing "greenstone schist." Creighton mine

x 8/10




Fig. 7




Fig. 8




Fig. 9



PLATE II

PHOTOMICROGRAPHS OF THIN SECTIONS OF LEAN ORE FROM THE STOBIE MINE, SUDBURY

FIG. 11

Typical norite with euhedral magnetite [black], hypersthene [gray], plagioclase [white].

Magnification x 36 diameters.

FIG. 12

Ore-minerals; magnetite, pyrrhotite [black], surrounding and replacing hypersthene [gray]. A veinlet of chlorite [ch] cuts magnetite. In the upper portion of the photograph the hypersthene is fresh, and in the lower part it is altered; and yet the ore shows the same relation to the silicates in the fresh and altered portions.

x 30

FIG. 13

A veinlet of chlorite [ch], cutting the sulfids, chalcopyrite and pyrrhotite [black], and the silicates.

x 36

FIG. 14

Tremolite [tr] crystals cutting—not residual in—chalcopyrite [black], and projecting from altered hypersthene [gray].

x 119




Fig. 11




Fig. 12




Fig. 13




Fig. 14

PLATE III

PHOTOMICROGRAPHS OF POLISHED SECTIONS OF ORES FROM THE STOBIE MINE, SUDBURY

FIG. 15

Lean ore. Chlorite veinlets cutting ore-minerals; magnetite, pyrrhotite, chalcopyrite [light], and the silicates.

x 11

FIG. 16

Plagioclase [dark], uralite [light gray], and sulfids [white]. Figs. 17 and 18 are higher magnifications from the same field.

x 10

FIG. 17

Veinlet and crystals of tremolite [gray] cutting pyrrhotite.

x 270

FIG. 18

Pyrrhotite [*p*] and pentlandite [*pn*]₁ cut by irregular veinlets of secondary silicates [*ss*]. A second generation of pentlandite (?) [*pn*]₂ grows out in tufts from these veinlets. The secondary migration of ore-minerals is therefore insignificant in amount.

Aggregates of tremolite crystals on the right of the photograph.

x 165




Fig. 15




Fig. 16



Fig. 17




Fig. 18

PLATE IV
PHOTOMICROGRAPHS OF ORES FROM SUDBURY

FIG. 19

Polished section of rich ore from the Stobie mine. Irregular areas and veinlets of chalcopyrite [white] replacing garnet [g] and other silicates.

x 16

FIG. 20

Copper Cliff mine. Thin section of ore in coarse granitic material. Ore-minerals [black areas], magnetite and pyrrhotite, cutting and replacing biotite.

x 38

FIG. 21

Polished section, Creighton mine. Crystallographic intergrowth of magnetite and ilmenite [parallel dark lines]. Specimen is not etched, but contrast is brought out by high polish.

x 790

FIG. 22

Polished section, Creighton mine. Typical rich ore showing pyrrhotite [p] with veins of pentlandite [pn] and residual magnetite [m] and silicates [s].

x 14




Fig. 19



Fig. 20




Fig. 21




Fig. 22

PLATE V

PHOTOMICROGRAPHS OF THIN SECTIONS OF ORE FROM SUDBURY

FIG. 23

Ore from the Creighton mine. Large, irregular ore masses [black] are chalcopyrite and pyrrhotite. Small subhedral black spots are chalcopyrite, probably a replacement of magnetite, as they contain residual specks of the latter. Ore-minerals cut through the silicates and replace them. Veinlets at the bottom of photograph are chalcopyrite.

x 28

FIG. 25

Creighton mine. Sulfid veinlets, chalcopyrite and pyrrhotite, which extend out from a mass of sulfids [black], cut silicates [hornblende and plagioclase].

x 27

FIG. 24

Creighton mine. Chalcopyrite veinlets [black] cut microcline, quartz, and biotite [dark gray], which are free from alteration products.

x 66

FIG. 26

Evans mine. Chalcopyrite veinlet cuts hornblende [*ho*], biotite [*bi*], and plagioclase [white]. Displacement, shown by hornblende crystal, has taken place along this veinlet. Beyond the field of the photograph chlorite cuts across the veinlet. [*ap*] apatite.

x 158



Fig. 23



Fig. 24




Fig. 25




Fig. 26

PLATE VI

PHOTOMICROGRAPHS OF POLISHED SECTIONS OF "ACID" ORE-BEARING ROCK FROM THE CREIGHTON MINE, SUDBURY

FIG. 27

Ore masses, pyrrhotite and chalcopyrite, extend out into fine veinlets, chiefly chalcopyrite. Large masses replace the silicates. The veinlets cut the subgraphic intergrowth of quartz [*q*] and feldspar [*f*]

x 10

FIG. 28

Sulfids, pyrrhotite [*p*] and chalcopyrite [*cp*] cut magnetite [*m*] and silicates in anastomosing veinlets.

x 15

FIG. 29

Chalcopyrite [*cp*] extending from the main mass into veinlets as one generation of ore. No evidence of "rearrangement" or second generation of ore in veinlets. A mass of pentlandite [*pn*] enclosed in chalcopyrite.

x 99

FIG. 30

Sharp veinlets of chalcopyrite cut the graphic intergrowth of quartz [dark] and feldspar [light].

x 260



Fig. 27




Fig. 28




Fig. 29




Fig. 30

PLATE VII

PHOTOMICROGRAPHS OF POLISHED SECTIONS OF "ACID" ORE-BEARING ROCK FROM THE CREIGHTON MINE, SUDBURY

FIG. 31

Selective replacement by sulfids [white] of feldspar [light gray] in the feldspar-quartz intergrowth. Quartz is dark gray.

x 18

FIG. 32

Sulfids, pyrrhotite [*p*] and chalcopyrite [*cp*] and veinlike masses of pentlandite [*pn*]. In the center of the field chalcopyrite preserves the structure of the graphic intergrowth by selective replacement of the feldspar.

x 18

FIG. 33

Graphic intergrowth of quartz and feldspar. A small irregular replacement veinlet cuts the fine graphic intergrowth above. Large-scale graphic intergrowth below preserved by selective replacement of feldspar by sulfids.

x 71

FIG. 34

An irregular area of magnetite replacing silicates.

x 140




Fig. 31




Fig. 32




Fig. 33




Fig. 34

PLATE VIII

PHOTOMICROGRAPHS OF POLISHED SECTIONS OF ORES FROM SUDBURY

FIG. 35

Massive ore from the Creighton mine showing typical vein-like areas of pentlandite $[pn]_1$ in pyrrhotite $[p]$. Contrast brought out by polishing, not by etching. Tiny brush-like crystals of a second generation of pentlandite $[pn]_2$ develop along veinlets and contacts. See Fig. 36.

x 18

FIG. 36

Massive ore from the Creighton mine. Second generation of pentlandite $[pn]$ along a veinlet of chalcopyrite $[cp]$, cutting pyrrhotite $[p]$. This rearrangement of ore-minerals is minor in amount, and can rarely be noted even under high powers. (Same specimen as shown in Fig. 35.)

x 670

FIG. 37

Massive ore from the Vermilion mine, showing pyrite veinlets [dark gray] cutting polydymite (?) $[po]$ and chalcopyrite $[cp]$. Note that pyrite is definitely later than the typical magmatic ore-minerals in all the occurrences figured.

x 10

FIG. 38

Ore from the Worthington mine. Pyrite [white] with a reticulate structure, cutting gangue [black] and sphalerite [gray].

x 58




Fig. 35



Fig. 36




Fig. 37




Fig. 38

PLATE IX

PHOTOMICROGRAPHS OF POLISHED SECTION OF ORE FROM THE ALEXO MINE, NORTHERN ONTARIO, CANADA

FIG. 39

Shows the general relations of the ore-minerals [large white areas] to the serpentinized silicates. The original ore-minerals, chiefly pyrrhotite and pentlandite with subordinate chalcopyrite and magnetite, surround and embay the serpentine pseudomorphs after olivine.

A minor amount of the second generation of chalcopyrite has been concentrated by serpentinization within the silicates.

x 17

FIG. 40

Enlargement of a part of the field in the upper left-hand portion of Fig. 39. Shows serpentine pseudomorphous after olivine, surrounded by magmatic sulfids, the contacts of which are not modified by serpentinization. The latter, however, affects a secondary concentration of chalcopyrite developed as a border between two different types of serpentine.

x 70




Fig. 39




Fig. 40

PLATE X

PHOTOMICROGRAPHS OF ORE FROM THE ALEXO MINE, FROM THE SAME
POLISHED SECTION FIGURED IN PLATE IX

FIG. 41

Pyrrhotite [*p*], pentlandite [*pn*],
chalcopyrite [*cp*]₁, and magnetite
[*m*] cut by veinlets of serpentine.

x 44

FIG. 42

First [*pn*]₁ and second [*pn*]₂ gen-
erations of pentlandite in pyrrho-
tite [*p*]. The two generations can
be distinguished by a slight differ-
ence in relief and color. Serpentine
veinlets are later than the first gen-
eration pentlandite and closely con-
nected with the second generation.
Pentlandite and serpentine tend to
develop along the crystallographic
lines of the pyrrhotite.

x 175

FIG. 43

Pyrrhotite [*p*] and pentlandite
[*pn*] of the first generation, cut by
a serpentine veinlet with a center of
magnetite [*m*], along which pent-
landite of the second generation
[*pn*]₂ develops.

x 610

FIG. 44

Showing later generations of chal-
copyrite [*cp*]₂ connected with dif-
ferent stages of serpentinization. A
large mass of pyrrhotite [*p*] is
shown near the top of the photo-
graph. Small dots in center may be
a third generation of chalcopyrite.

x 183



Fig. 41



Fig. 42



Fig. 43



Fig. 44

PLATE XI

PHOTOMICROGRAPHS OF THIN SECTION OF PYRRHOTITE-BEARING PYROXENITE FROM THE GOLDEN CURRY MINE, ELKHORN, MONTANA

FIG. 45

Sulfids [black], chiefly pyrrhotite with some chalcopyrite, surrounding and penetrating augite. White needles in the sulfids in the center of photograph are tremolite and darker patches to the right are hornblende.

x 22

FIG. 46

Sulfids replacing hornblende along cleavages.

[The white streak (lower center) is a crack in the slide.]

x 130

FIG. 47

Sulfids surrounding and penetrating pyroxene. A hornblende crystal [ho] in the center is bordered by tremolite [tr] needles which cut the ore-minerals.

x 90




Fig. 45




Fig. 46




Fig. 47

PLATE XII

PHOTOMICROGRAPHS OF POLISHED SECTIONS FROM EVJE, NORWAY, AND
SOHLAND, GERMANY

FIG. 48

Ore from the Flaad mine, Evje, Norway.

Pyrite [*py*] veinlet and detached crystals in pyrrhotite [*p*] and magnetite [*m*]. The veinlet is connected with alteration products. See fig. 49 for the relation of pyrite crystals to pyrite veinlets.

x 12

FIG. 49

Ore from the Flaad mine, Evje, Norway.

Pyrrhotite [*p*] cut by chalcopyrite [*cp*] which in turn is cut by pyrite [*py*] veinlets. The latter connect with subhedral and euhedral crystals. The euhedral crystals are shown in fig. 48.

x 300

FIG. 50

Ore from Sohland, Saxony.

Shows general relations. The ore-minerals are pyrrhotite [*p*], chalcopyrite [*cp*], and pentlandite [*pn*]. Uralite needles [*u*] penetrate into the sulfids. Other silicates [*s*] are hornblende and alteration products. A good example of extensive post-mineral alteration.

x 12

FIG. 51

Ore from Sohland, Saxony.

Uralite [*u*] needles cutting pyrrhotite [*p*] and pentlandite [*pn*]. The uralite (or tremclite) develops at the ends of the hornblende crystals.

x 116



Fig. 48




Fig. 49




Fig. 50




Fig. 51

PLATE XIII

PHOTOMICROGRAPHS OF SECTIONS OF ORE-BEARING HYPERSTHENITE FROM NABABEEP MINE, OOKIEP, SOUTH AFRICA

FIG. 52

Thin section showing ore-minerals [black], magnetite and a little bornite, cutting and surrounding hypersthene.

x 74

FIG. 53

Thin section of hypersthene showing euhedral crystals of magnetite within the hypersthene and anhedral ore-minerals, chiefly magnetite, connected with the "acid extract" [white]. The "acid extract" is plagioclase.

x 59

FIG. 54

Polished section of ore-bearing hypersthene. The ore-minerals are magnetite [*m*] and bornite [*b*]. The general relations suggest that magnetite is partially replaced by bornite.

x 10

FIG. 55

Thin section showing anthophyllite [*an*] and talc (?) [*ta*] in pyrrhotite [black]. The sharp-pointed anthophyllite needles cut and are later than the ore-minerals.

x 435



Fig. 52




Fig. 53




Fig. 54




Fig. 55

PLATE XIV

PHOTOMICROGRAPHS OF THIN SECTIONS OF ORE-BEARING NORITE FROM THE TWEEFONTEIN MINE, OOKIEP, SOUTH AFRICA

FIG. 56

Hypersthene [dark gray] and plagioclase [light gray] surrounded and cut by ore-minerals, magnetite and a little chalcopyrite. Euhedral magnetite [designated by arrow] is developed within the silicates. Silicates are free from any kind of alteration products.

x 17

FIG. 57

A higher magnification of the lower portion of fig. 56. Magnetite is anhedral along the boundaries of the silicates and euhedral [see arrow] within the plagioclase.

x 49

FIG. 58

Hypersthene [gray] and plagioclase [white] surrounded and replaced by ore-minerals. Anhedral areas are chalcopyrite and bornite. Subhedral crystals within the hypersthene are magnetite.

x 11

FIG. 59

A higher magnification of the plagioclase in the left side of fig. 58. The ore-minerals are "eating in" along albite twinning lamellae.

x 23




Fig. 56



Fig. 57



Fig. 58



Fig. 59

PLATE XV

PHOTOMICROGRAPHS OF A POLISHED SECTION OF ORE-BEARING NORITE, TWEEFONTEIN MINE, OOKIEP, SOUTH AFRICA

FIG. 60

The ore-minerals shown in the slide are bornite [*b*], chalcopyrite [*cp*], magnetite [*m*], lined with ilmenite. (Hematite is shown in fig. 61.) Silicates are plagioclase [*pl*], hypersthene [*hy*], biotite [*bi*], and apatite [*ap*]. Note especially magnetite cutting hypersthene in the upper portion of the photograph, and bornite and chalcopyrite penetrating biotite on the left.

x 9

FIG. 62

Photograph from the same section as shown in figs. 60 and 61. Chalcopyrite [*cp*]₂, and bornite [*b*] cut by a veinlet of anthophyllite [black needles], along which is developed a second generation of chalcopyrite (shown faintly at [*cp*]₂).

x 150

FIG. 61

Photograph from the same polished section as shown in fig. 60, showing hematite [*h*], and magnetite [*m*] intergrown with ilmenite lamellae.

x 18

FIG. 63

A higher magnification of the veinlet shown in fig. 62. Bornite [gray], anthophyllite [black], and chalcopyrite of the second generation [white].

x 920




Fig. 60




Fig. 61




Fig. 62



Fig. 63

PLATE XVI

PHOTOMICROGRAPHS OF SECTIONS OF ORE-BEARING MICA DIORITE, OOKIEP EAST MINE, OOKIEP, SOUTH AFRICA

FIG. 64

Polished section showing pyrrhotite [*p*] residual in chalcopyrite [*cp*] and sulfid veinlets in the silicates.
x 10

FIG. 65

Thin section. The ore-minerals [black] are pyrrhotite and chalcopyrite. The primary silicate is plagioclase [gray]. The secondary silicates cut the ore-minerals and plagioclase as veinlets in the lower portion of photograph, and are distributed as specks in the sulfids. The alteration is subsequent to ore-formation.
x 18

FIG. 66

From the same thin section as fig. 65. The veinlets of sulfids are not a later generation of ore, nor are they due to rearrangements. This fact is shown in fig. 69.
x 45

FIG. 67

A group of subhedral magnetite crystals [*m*], a few of which are partially replaced by pyrrhotite [*p*] [at the bottom of the photograph].
x 51




Fig. 64




Fig. 65




Fig. 66



Fig. 67

PLATE XVII

PHOTOMICROGRAPHS OF POLISHED SECTIONS OF ORE-BEARING MICA DIORITE, OOKIEP EAST MINE, OOKIEP, SOUTH AFRICA

FIG. 68

Pyrrhotite [*p*] and chalcopyrite [*cp*] cutting the feldspars in sharp veinlets and penetrating the cleavage planes of the biotite [top of photograph].

x 10

FIG. 70

A biotite-rich segregation in the diorite, with the sulfids, chalcopyrite [*cp*] and pyrrhotite [*p*], penetrating the cleavages of the biotite.

x 11

FIG. 69

The chalcopyrite veinlet on the right is cut sharply by chloritic alteration products, and therefore the veinlets are not connected with the rock alteration, but antedate the latter. On the left a biotite crystal [*bi*] is penetrated by chalcopyrite along cleavage planes.

x 153

FIG. 71

Brush-like pentlandite (?) [*pn*], replacing pyrrhotite [*p*] and chalcopyrite [*cp*]. It is probably of a late generation, like that shown in figs. 18, 35, 36, 42 and 43.

x 620




Fig. 68



Fig. 69




Fig. 70



Fig. 71

PLATE XVIII

PHOTOMICROGRAPHS OF THIN SECTIONS OF NORITE-DIORITE FROM THE ENGELS MINE, PLUMAS COUNTY, CALIFORNIA

FIG. 72

Unaltered norite-diorite showing the general relations of the ore-minerals [black] to the silicates (hypersthene [dark gray] and feldspar [light gray]). The ore-minerals are chiefly magnetite and hematite. The magnetite shows all gradations in form from euhedral to anhedral outlines.

x 22

FIG. 73

A higher magnification of a spot in the upper right-hand corner of fig. 72. Euhedral magnetite occurs within the silicates, anhedral along the boundaries. One area [indicated by arrow] shows an euhedral termination penetrating a silicate crystal, and the remaining portion is anhedral.

x 143

FIG. 74

A higher magnification of another spot in the upper right corner of fig. 72. Hook-shaped anhedral magnetite and hematite [black] penetrating and replacing the silicates, especially biotite [bi].

x 195

FIG. 75

Hornblende [ho] rim surrounding pyroxenes [px] (diopside and hypersthene). This phenomenon is often observed in magmatic deposits.

x 40




Fig. 72



Fig. 73



Fig. 74




Fig. 75

PLATE XIX

PHOTOMICROGRAPHS OF SECTIONS OF ROCKS AND ORES FROM THE ENGELS MINE, PLUMAS COUNTY, CALIFORNIA

FIG. 76

Thin section of norite-diorite showing diopside [*di*] with rim of hornblende [*ho*]. Euhedral crystals of magnetite [black] within the diopside, and anhedral crystals of magnetite and hematite [black] along its borders.

x 63

FIG. 77

Thin section of norite-diorite showing silicates [light gray and white] and ore-minerals [black], magnetite and hematite, cut by veinlets of chlorite (shown by arrow), which proves that the alteration is post-mineral.

x 36

FIG. 78

Thin section of ore-bearing granodiorite containing tourmaline [dark gray], sericitized feldspars [light gray], and ore-minerals [black], bornite and chalcocite. The feldspars and tourmaline are replaced by the ore-minerals and they in turn by sericite laths. (One is indicated by arrow in the lower right-hand corner.) This section shows local development of sericite. Most specimens, however, are not affected by sericitization.

x 23

FIG. 79

Polished section of the ore shown in fig. 78. Note the irregular hook-shaped bornite [*b*] with narrow rim of supergene chalcocite [*cc*]₂. The bornite surrounds and replaces the silicates.

x 184



Fig. 76



Fig. 77




Fig. 78



Fig. 79

PLATE XX

PHOTOMICROGRAPHS OF POLISHED SECTIONS OF ORES FROM THE ENGELS MINE, PLUMAS COUNTY, CALIFORNIA, SHOWING LOCAL REARRANGEMENT AND COPPER ENRICHMENT SUB- SEQUENT TO THE MAGMATIC STAGE

FIG. 80

The photograph shows the following post-magmatic alterations of bornite [b]: First the development of the so-called graphic intergrowth of bornite [b] and chalcocite [cc]₁, and later the development of rims of chalcocite [cc]₂ around the margins of the bornite areas. The first generation of chalcocite is probably hypogene, the second generation probably supergene.

x 125

FIG. 82

An area of bornite [b] has been replaced by covellite [cv] and chalcocite [cc]₁, and later all of these have been penetrated by chlorite [ch] laths. This alteration is in part along crystallographic directions of the bornite.

x 142

FIG. 81

Anhedron of bornite [b] altering to covellite [cv] and chalcopyrite [cp]₂ of the second generation. The latter develops along crystallographic directions of the bornite. This alteration is probably supergene.

x 542

FIG. 83

An area of bornite [b] has been replaced by chlorite [ch] laths and by quartz [q] veinlets. Chalcocite [cc]₂ of the second generation has developed along the margin of the bornite, along the chlorite-bornite contacts, and along veinlets.

x 142




Fig. 80



Fig. 81



Fig. 82




Fig. 83

- ACCELERATION IN THE DEVELOPMENT OF THE FOSSIL CEPHALOPODA.** James Perrin Smith, Professor of Paleontology. 30 pp., 15 plates. 1914. Price, 75c.
- A MORPHOLOGICAL STUDY OF SOME MEMBERS OF THE GENUS PALLAVINIA.** Douglas Houghton Campbell, Professor of Botany, and Florence Williams. 44 pp., 23 text figures. 1914. Price, 50c.
- THE EVOLUTION OF BRAZIL COMPARED WITH THAT OF SPANISH AND ANGLO-SAXON AMERICA.** Manuel de Oliveira Lima, Minister of Brazil to Belgium. 160 pp. 1914. Price, \$1.00.
- THE HEMOLYMPH NODES OF THE SHEEP.** Arthur William Meyer, Professor of Anatomy. 74 pp., 1 plate, 4 colored plates. 1914. Price, \$1.00.
- AN INTRODUCTION TO THE STUDY OF THE ENDOCRINE GLANDS AND INTERNAL SECRETIONS.** (Lane Medical Lectures.) Sir Edward Schäfer, Regius Professor of Physiology in the University of Edinburgh. 94 pp. 1914. Price, 75c.
- THE PRONOUN OF ADDRESS IN ENGLISH LITERATURE OF THE THIRTEENTH CENTURY.** Arthur Garfield Kennedy, Instructor in English Philology. 91 pp. 1915. Price, \$1.00.
- THE ANOPLURA AND MALLOPHAGA OF NORTH AMERICAN MAMMALS.** Vernon Lyman Kellogg, Professor of Entomology, and Gordon Floyd Ferris. 74 pp., 18 text figures, 8 plates. 1915. Price, 75c.
- THE FLÜGEL MEMORIAL VOLUME.** Papers by Ewald Flügel, late Professor of English Philology, his Colleagues and Students. 232 pp. 1916. Price, \$1.50.
- THE SESAMOID ARTICULAR: A BONE IN THE MANDIBLE OF FISHES.** Edwin Chapin Starks, Assistant Professor of Zoology. 40 pp., 15 text figures. 1916. Price, 50c.
- A STUDY OF GERMAN VERBS COMPOUNDED WITH AUS, EIN, ETC. AS CONTRASTED WITH THOSE COMPOUNDED WITH HERAUS, HINAUS, HEREIN, HINEIN, ETC.** Charles Reining. 142 pp. 1916. Price, \$1.00.
- THE PATHOLOGY OF NEPHRITIS.** William Ophüls, Professor of Pathology. 103 pp., 64 text figures and plates. 1916. Price, \$1.00.
- BONE AND JOINT STUDIES, I.** Leonard W. Ely, Associate Professor of Surgery, and John Francis Cowan, Assistant Professor of Surgery. 139 pp., 41 text figures and plates. 1916. Price, \$1.00.
- A STUDY OF THE MAGMATIC SULFID ORES.** C. F. Tolman, Jr., Associate Professor of Economic Geology, and Austin F. Rogers, Associate Professor of Mineralogy and Petrography. 76 pp., 7 text figures, 20 plates. 1916. Price, \$1.00.

To avoid fine, this book should be returned on
or before the date last stamped below

SDM-8-40

NO 1 46

FEB 26 1993

FEB 11 1994

553.16 .T652

C.1

... A study of the magmatic su

Stanford University Libraries



3 6105 032 254 026

553.
T652

31776